

Older adults who are frail more likely to experience delirium after surgery

January 26 2018

Older adults who are frail are twice as likely to experience delirium following elective surgery than those of an older age, a new study suggests.

Researchers at St. Michael's Hospital found that a history of [delirium](#), frailty and cognitive impairment are the risk factors most strongly associated with developing postoperative delirium, according to the systematic review, published online today in the *Journal of General Internal Medicine*.

Previous research by the authors found that frailty and [cognitive impairment](#) before [surgery](#) are associated with developing complications after surgery, but age is not.

Other risk factors that are associated with developing postoperative delirium include smoking and the use of psychotropic medications, according to the authors.

"Chronological age from your birth date is not always an accurate assessment of how you've aged over your lifetime," said Dr. Jennifer Watt, a geriatric medicine fellow at St. Michael's and lead author of the review. "This study highlights how common delirium is among [older adults](#) undergoing elective surgery, and the importance of geriatric syndromes, including frailty, in identifying older [adults](#) who may be at risk."

The review examined 41 existing studies including more than 9,000 patients 60 years and older and reporting on [postoperative delirium](#) following elective surgery.

The researchers found that one in six older adults experience delirium following elective surgery. They also found that patients who had caregiver support are also 30 per cent less likely to experience delirium following elective surgery than those who did not.

Caregiver support was defined in the studies as the patient being married or having a higher number of visitors than average, according to the review. The studies included in the review did not specify whether caregiver support came before surgery, after surgery, or both, according to the authors.

"Previous research has shown that when families delivered a non-pharmacological intervention, such as helping to mobilize or orient the patient, the patient was less likely to experience delirium following hip surgery," said Dr. Watt. "What we're seeing in this review may be a result of the fact that they're doing some of these things for their loved ones already."

Patients who developed delirium following elective surgery are also at an increased risk of developing other adverse postsurgical outcomes, and are more likely to be discharged to another hospital, a long-term care facility, or die following surgery, according to the review.

Although older adults are assessed for cardiovascular and respiratory risk before an elective surgery, often not enough consideration is given to risk factors that are more common in older adults, including delirium, the authors said.

"Postoperative delirium is a common, yet preventable, complication

experienced by older adults undergoing [elective surgery](#)," said Dr. Watt. "Understanding delirium [risk factors](#) may help clinicians, patients, and caregivers to target interventions aimed at lessening its burden."

Provided by St. Michael's Hospital

Citation: Older adults who are frail more likely to experience delirium after surgery (2018, January 26) retrieved 2 May 2024 from <https://medicalxpress.com/news/2018-01-older-adults-frail-delirium-surgery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--