

Oxysterols guide gut immune cells and are involved in inflammatory bowel disease

January 16 2018

Researchers at Karolinska Institutet in Sweden report that cholesterol metabolites cause specific immune cells in the large intestine to move, which lies behind the formation of the immune system's important lymphoid tissue in the intestine. The study, published in the journal *Immunity*, paves the way for a new possible treatment for patients with inflammatory bowel disease.

Inflammatory bowel disease is a common and severe disease that has no cure. Normally, the immune cells of the gut accumulate and communicate with each other in the <u>lymphoid tissue</u>. But, for those who have autoimmune <u>inflammatory bowel disease</u>, such as <u>ulcerative colitis</u>, the lymphoid tissue can react to the body's own cells and cause chronic inflammation.

Hitherto, it has not been known which signals control the formation of the lymphoid tissue in the <u>large intestine</u>. But, now scientists at Karolinska Institutet have discovered that cholesterol metabolites, called oxysterols, stimulate specific immune cells in the large intestine, called <u>innate lymphoid cells</u> (ILCs), to form lymphoid tissue in both health and in disease.

"We have discovered that ILCs use a surface protein called GPR183 to sense the oxysterols, which attracts ILCs to specific sites in the large intestine where lymphoid tissue is formed," says Tim Willinger, researcher at Karolinska Institutet's Department of Medicine in Huddinge, who has led the study.



"Our findings are relevant to humans because inflammatory lymphoid tissue contributes to tissue damage in inflammatory bowel disease. We have discovered that patients with ulcerative colitis have higher levels of oxysterol-producing enzymes than healthy controls. The results indicate that oxysterols and GPR183 are involved in ulcerative colitis," he says.

One way to treat inflammatory bowel disease could, therefore, be to attempt to inhibit GPR183 or oxysterol production.

"This treatment option is attractive, because GPR183 belongs to a group of molecules that are excellent targets for drug treatment," says Dr Willinger.

More information: "Oxysterol sensing through the receptor GPR183 promotes the lymphoid tissue-inducing function of innate lymphoid cells and colonic inflammation". Johanna Emgård, Hana Kammoun, Bethania García-Cassani, Julie Chesné, Sara M. Parigi, Jean-Marie Jacob, Hung-Wei Cheng, Elza Evren, Srustidhar Das, Paulo Czarnewski, Natalie Sleiers, Felipe Melo-Gonzalez, Egle Kvedaraite, Mattias Svensson, Elke Scandella, Matthew R. Hepworth, Samuel Huber, Burkhard Ludewig, Lucie Peduto, Eduardo J. Villablanca, Henrique Veiga-Fernandes, João P. Pereira, Richard A. Flavell, Tim Willinger. *Immunity*, online 16 January 2018.

Provided by Karolinska Institutet

Citation: Oxysterols guide gut immune cells and are involved in inflammatory bowel disease (2018, January 16) retrieved 4 May 2024 from https://medicalxpress.com/news/2018-01-oxysterols-gut-immune-cells-involved.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.