

Prolonged sedentary periods associated with fat around internal organs

January 9 2018



Credit: CC0 Public Domain

A team of researchers from the NIHR Leicester Biomedical Research Centre, UK – a partnership between Leicester's Hospitals, the University of Leicester and Loughborough University—has found new evidence to

suggest that longer periods of sedentary time (defined as any sitting/reclining activity with low energy expenditure) are more strongly associated with the amount of fat deposited around internal organs.

The study, which was published in the journal, *Obesity*, took 124 participants at high risk of type 2 diabetes and measured the length of time they spent sedentary over a period of seven days using accelerometers fitted to their waist. The research team also scanned the participants using magnetic resonance imaging (MRI) equipment to accurately measure the amount of fat in the liver, inner (visceral) and outer (subcutaneous) fat layers, and total abdominal fat.

Even when accounting for age, ethnicity and [physical activity](#) levels, the study team found that the longer a person remained sedentary during the day, the higher the levels of liver fat, inner (visceral) fat and total abdominal fat. The link was even stronger for participants who did not meet Public Health England's recommendation of 150 minutes of [moderate intensity](#) physical [activity](#) per week.

Dr. Joe Henson, Research Associate at the University of Leicester, led the study. He said: "We know that spending long periods of time sedentary is unhealthy and a risk factor for chronic illnesses, such as type 2 diabetes and heart disease. Likewise, the amount of fat deposited around our internal organs may also predispose us to these diseases.

"Using MRI techniques and physical activity monitors we have shown that the more time spent sedentary, the stronger the association with higher levels of internal and abdominal fat. This was particularly so if the long periods of sedentary behaviour were uninterrupted. Our findings also show that reaching the UK government's target of 150 minutes of moderate intensity physical activity may offer some protection against the harmful effects of prolonged sedentary time."

Professor Melanie Davies, Director of the NIHR Leicester Biomedical Research Centre, UK, Professor of Diabetes Medicine at the University of Leicester and a co-author of the study, said: "Lack of physical activity and being overweight are two risk factors associated with type 2 diabetes. However, the effects of prolonged sedentary time and whether physical activity can play a mediating role by reducing fat deposits on [internal organs](#) remain unclear. This research starts to shed a light on any connections between the two by using MRI to measure the distribution of fat in an individual's body and analysing that in relation to their activity levels. The next step would be to examine the impact of regularly breaking up prolonged [sedentary time](#) upon internal fat levels."

More information: Joseph Henson et al. Sedentary Time and MRI-Derived Measures of Adiposity in Active Versus Inactive Individuals, *Obesity* (2017). [DOI: 10.1002/oby.22034](https://doi.org/10.1002/oby.22034)

Provided by University of Leicester

Citation: Prolonged sedentary periods associated with fat around internal organs (2018, January 9) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-01-prolonged-sedentary-periods-fat-internal.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--