

Researchers identify protein involved in cocaine addiction

January 16 2018



A pile of cocaine hydrochloride. Credit: DEA Drug Enforcement Agency, public domain

Mount Sinai researchers have identified a protein produced by the immune system—granulocyte-colony stimulating factor (G-CSF)—that could be responsible for the development of cocaine addiction.

A study showed that G-CSF can alter a mouse's desire for cocaine, but not for other rewards. This effect is modulated by a brain region that plays a central role in [reward processing](#) and addiction. If applicable to humans, these findings represent a potential therapeutic approach to decrease a cocaine addict's motivation to seek the drug without introducing a potential new substance for abuse.

The results of the study will be published online in *Nature Communications*.

Previous research has demonstrated a link between cocaine use and the immune system in humans and animals, with addicts showing altered immune responses to drugs and drug cues.

In this study, the research team identified G-CSF—a cytokine produced by immune cells which was expressed at higher levels in both the blood and brain in mice that were treated with repeated doses of cocaine. Injecting G-CSF into the [nucleus accumbens](#), a brain region associated with reward, causes mice to take more cocaine, but does not change their motivation to consume a more natural reward, sugar water. Conversely, injecting an antibody that neutralizes G-CSF in the nucleus accumbens can reduce the mouse's motivation to take cocaine. Taken together, the results from this study suggest that manipulating G-CSF in the [reward center](#) of the brain changes the biochemical signals that push animals to take cocaine.

"The results of this study are exciting because outside of 12-step programs and psychotherapy, no medication-assisted therapy exists to treat [cocaine](#) addiction," said the study's senior author, Drew Kiraly, MD, PhD, Assistant Professor of Psychiatry at Icahn School of Medicine at Mount Sinai. "Drugs that manipulate G-CSF already exist as FDA-approved medications. Once we clarify how G-CSF signaling can best be targeted to reduce addiction-like behaviors, there is a high

possibility that treatments targeting G-CSF could be translated into clinical trials and treatments for patients."

More information: Granulocyte-colony stimulating factor controls neural and behavioral plasticity in response to cocaine, *Nature Communications* (2018).

[nature.com/articles/doi:10.1038/s41467-017-01881-x](https://doi.org/10.1038/s41467-017-01881-x)

Provided by The Mount Sinai Hospital

Citation: Researchers identify protein involved in cocaine addiction (2018, January 16) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2018-01-protein-involved-cocaine-addiction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.