

Psychiatric medications are not overprescribed for kids, finds study

January 29 2018



Credit: CC0 Public Domain

A new study at Columbia University Irving Medical Center (CUIMC) challenges the popular notion that psychiatric medications are overprescribed in children and adolescents in the U.S. When the

researchers compared prescribing rates with prevalence rates for the most common psychiatric disorders in children, they discovered that some of these medications may be underprescribed.

The findings were published online today in the *Journal of Child and Adolescent Psychopharmacology*.

"Over the last several years, there has been widespread public and professional concern over reports that [psychiatric medications](#) are being overprescribed to [children](#) and adolescents in the United States," Ryan Sultan, MD, a child psychiatrist and researcher at CUIMC who led the study. "We were interested in better understanding this concern."

How the study was done

Using data from a national prescription database, the researchers looked at annual [prescriptions](#) for three psychiatric drug classes-stimulants, antidepressants, and antipsychotics-for 6.3 million children between the ages of 3 and 24 years. They then compared prescribing patterns with known prevalence rates of attention deficit-hyperactivity disorder (ADHD), anxiety disorders, and depression between young children (3 to 5 years), older children (6 to 12 years), adolescents (13 to 18 years), and young adults (19 to 24 years). This is the first national study to analyze prescription rates for these three types of psychiatric medications in youth.

Annually, an estimated 1 in 8 U.S. teenagers has a depressive episode, and roughly 1 in 12 children have symptoms of ADHD. During the year studied, fewer than 1 in 30 teenagers received a prescription for antidepressants, and only 1 in 20 got a prescription for stimulants.

"Our results show that, at a population level, prescriptions of stimulants and antidepressant medications for children and adolescents do not

appear to be prescribed at rates higher than the known rates for psychiatric conditions they are designed to treat," said Dr. Sultan. "These findings are inconsistent with the perception that children and adolescents are being overprescribed."

Overall psychiatric drug prescription patterns in children and adolescents Children in the youngest group accounted for the smallest number (0.8 percent) of prescriptions for any psychiatric drug. Adolescents accounted for the highest number (7.7 percent).

The number of prescriptions for stimulants was highest in older children (4.6 percent), with males accounting for more of these prescriptions than females. Antidepressant prescriptions increased with age and was highest for young adults (4.8 percent), particularly for females. Antipsychotic prescriptions peaked during adolescence (1.2 percent) and were prescribed slightly more often for males in this age group.

"The study also showed that, among young people in the United States, the patterns of prescriptions for antidepressants and stimulants are broadly consistent with the typical ages associated with the onsets of common mental disorders, said Mark Olfson, MD, professor of psychiatry at CUIMC and senior author of the paper. "However, the situation with antipsychotic medications is less clear cut. Given clinical uncertainty over their appropriate indications, it is unclear whether their annual use rates, which ranged from 0.1 percent in younger children to 1 percent in adolescents, are above or below the rates of the psychiatric disorders they aim to treat."

"These results provide some reassurance to those who are concerned about the overprescribing of psychiatric medications to children and teenagers," said Dr. Sultan. "Improving access to child psychiatrists through consultation services and collaborative care models may help address potential undertreatment while also reducing the risk of

prescribing medications before other treatments have been tried. Acknowledging that prescription patterns may have changed since the data was collected, it is important to continue to assess the pattern and distribution of psychiatric medications to children and adolescents in the U.S."

More information: Ryan S. Sultan et al. National Patterns of Commonly Prescribed Psychotropic Medications to Young People, *Journal of Child and Adolescent Psychopharmacology* (2018). [DOI: 10.1089/cap.2017.0077](https://doi.org/10.1089/cap.2017.0077)

Provided by Columbia University Medical Center

Citation: Psychiatric medications are not overprescribed for kids, finds study (2018, January 29) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-01-psychiatric-medications-overprescribed-kids.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--