

Sex education doesn't reflect real-life realities of lesbian and bisexual girls

January 10 2018



Michele Ybarra, Center for Innovative Public Health Research (CiPHR). Credit: Center for Innovative Public Health Research

Most lesbian and bisexual girls don't know they can get sexually transmitted infections (STIs) from other girls, because sex education is

mostly designed for their straight peers. This knowledge gap could be placing them at increased risk for getting STIs.

That's one of the conclusions of a new study led by researchers at the Centre for Innovative Public Health Research, a non-profit research group based in California, conducted in collaboration with researchers at the University of British Columbia and the City University of New York.

Previous UBC studies show that [lesbian](#) and bisexual [girls](#) face higher risks for teenage pregnancy than straight girls, says study co-author and UBC youth health researcher Jennifer Wolowic. In this new study, the researchers wanted to examine what lesbian and bisexual girls know about safe sex and their STI risks, especially when having sex with other girls.

"What surprised us was their overall lack of knowledge when it came to safe sex practices with female partners," said Wolowic. "When we asked why, many told us they didn't find their sex ed programs-if they even had one-to be very informative. And even when they asked questions, the focus on heterosexual sex made them feel uncomfortable."

The researchers conducted online focus groups with 160 lesbian and bisexual teenage girls located across the U.S. They found that a lack of knowledge and concerns about loss of pleasure were two of the main reasons for their limited use of dental dams and other barriers.

"Participants told us, they 'literally had never heard of dental dams,' or thought STIs weren't a concern when having sex with girls. Of those who knew about protective barriers, many said using protection made sex awkward or less pleasurable, and so they left them out during sex," said Wolowic.

Many teens thought that getting tested for [sexually transmitted infections](#) was something important to do in relationships. At the same time, they thought they could trust a female partner to be "clean" to a greater extent than a male partner.

"Add to that a lack of information about where to get dental dams or how to make them, and it's easy to understand why barrier use is so low among lesbian and bisexual girls," says Michele Ybarra, the study's principal investigator. "They need to know that there are sexy ways to use barriers, that they can make dental dams out of condoms if needed, and that they can get STIs having sex with other girls."

This study also highlights the need for more inclusive sex ed, adds UBC nursing professor Elizabeth Saewyc, the paper's senior author who leads the Stigma and Resilience Among Vulnerable Youth Centre at UBC.

"Young people need accurate sexual health information, but [sex education](#) has traditionally focused on heterosexual sex," said Saewyc. "Our findings suggest we need to create more inclusive curriculum to help lesbian and bisexual girls have the knowledge they need to make healthy sexual decisions."

More information: Marion Doull et al, Why Girls Choose Not to Use Barriers to Prevent Sexually Transmitted Infection During Female-to-Female Sex, *Journal of Adolescent Health* (2017). [DOI: 10.1016/j.jadohealth.2017.10.005](#)

Provided by University of British Columbia

Citation: Sex education doesn't reflect real-life realities of lesbian and bisexual girls (2018, January 10) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-01-sex-doesnt->

[real-life-realities-lesbian.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.