


A 'touching sight': How babies' brains process touch builds foundations for learning

January 16 2018, by Kim Eckart



These images illustrate two views of the left hemisphere of the brain. Image A shows the location where infants in the experiment processed a touch to the hand; image B shows where, in the brain, they processed a touch to the foot. Credit: UW Institute for Learning & Brain Sciences

Touch is the first of the five senses to develop, yet scientists know far less about the baby's brain response to touch than to, say, the sight of

mom's face, or the sound of her voice.

Now, through the use of safe, new brain imaging techniques, University of Washington researchers provide one of the first looks inside the infant's brain to show where the sense of touch is processed—not just when a baby feels a touch to the [hand](#) or foot, but when the baby sees an adult's hand or foot being touched, as well.

The evidence of activity in the somatosensory cortex for both "felt touch" and "observed touch" shows that 7-month-old [infants](#) have already made a basic connection between "self" and "other," which researchers say lays the groundwork for imitating and learning from the behavior of other people, and for empathizing with them.

The findings by the UW Institute for Learning & Brain Sciences (I-LABS) are published this week in *Developmental Science*.

"Long before babies acquire spoken language, touch is a crucial channel of communication between caregivers and babies," said the study's primary author, Andrew Meltzoff, UW psychology professor and co-director of I-LABS. "Now we have the tools to see how the baby's body is represented in the baby's brain. This allows us to catch the first glimpse of a primitive sense of self that provides a building block for social learning."

Past studies investigated how infants' brains respond to touch, generally. The authors believe this is the first experiment to measure the specific networks of the brain where this processing occurs, and to illuminate how babies' brains respond to seeing another person being touched, in the absence of being touched themselves.

For the study, researchers used the I-LABS Magnetoencephalography (MEG) machine to capture images of brain activity in 7-month-old

infants as they were touched on the hand and foot, and as they watched videos of an adult hand and foot being touched.

Researchers were particularly interested in the brain's somatosensory cortex, a region generally described as a strip of tissue in the brain that runs between the ears, over the top of the head. It is in this region, in separate places and at different levels of strength, that the brain processes touch to different parts of the body. A touch to the hand, for example, is a stronger sensation - and is processed in a different location along the somatosensory cortex - than a touch to the foot.

In the first experiment, each infant was seated in the MEG to measure brain activity as they received light touches. A small, inflatable balloon-like device was placed on the top of the baby's hand, and when it expanded and contracted according to a computer-controlled timetable, it produced light taps on the baby's skin. The same procedure was followed for the top of the baby's foot.

The data showed that, when the hand was touched, the hand area of the somatosensory cortex was activated in all 14 infants tested; when the foot was touched, activation occurred in the foot area of the brains of all of the infants but one.

A different group of infants provided data for the "observed touch" experiment, in which they also were seated in the MEG but watched separate videos of an adult hand and an adult foot being touched by a small rod. Researchers discovered that the infants' own somatosensory cortex (the "touch center" in the baby brain) also became activated when the babies simply observed someone else being touched.

There was a weaker response to "observed touch" than to "felt touch," which was expected, Meltzoff said. The same is true of adults: A touch to your own hand is going to generate greater [brain activity](#) in the

[somatosensory cortex](#) than merely seeing the touch to someone else's hand.

The key, Meltzoff pointed out, is that the same part of the infant's brain registered both kinds of touch, indicating a baby's capacity for recognizing the similarity between their own body parts and those they see in other people.

This new evidence for shared neural regions processing touch to self and [touch](#) to others makes sense, Meltzoff said. As parents know, babies watch and imitate what adults do. Imitation is a powerful learning mechanism for infants, but in order to imitate, infants have to perceive how body parts correspond. In other words, they need to reproduce the same movement with the same part when they imitate what their parent is doing. Scientists have wondered how infants make this connection. "Before they have words for the [body parts](#), babies recognize that their hand is like your hand, and their foot is like your [foot](#). The neural body map helps connect [babies](#) to other people: The recognition that another person is 'like me' may be one of the baby's first social insights," Meltzoff explained.

With development, this "like-me" recognition eventually flowers into feeling empathy for someone else. If you see someone accidentally hit their thumb with a hammer, you rapidly, if perhaps imperceptibly, recoil by moving your hand. This is where a shared neural body map that connects self to other comes into play.

Further research could use the MEG to investigate how infants develop more sophisticated body awareness as they grow older, the paper notes.

"The idea of using [brain](#) science to study how and when humans first feel a sense of connectedness with others is important and fascinating," Meltzoff said. "We can now look under the hood and see what's

happening when a baby watches and connects to others. It's a touching sight."

More information: Andrew N. Meltzoff et al, Infant brain responses to felt and observed touch of hands and feet: an MEG study, *Developmental Science* (2018). [DOI: 10.1111/desc.12651](https://doi.org/10.1111/desc.12651)

Provided by University of Washington

Citation: A 'touching sight': How babies' brains process touch builds foundations for learning (2018, January 16) retrieved 20 March 2024 from <https://medicalxpress.com/news/2018-01-sight-babies-brains-foundations.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--