

Light activity measured with fitness tracker linked to lower mortality in older women

January 10 2018

Experts say that a lack of physical activity leads to age-related weakness and poor health in older adults. Official guidelines suggest that healthy older adults spend at least 2.5 hours every week doing moderate activity (such as brisk walking), or at least 1.25 hours per week doing vigorous exercise (such as jogging or running).

Unfortunately, many [older adults](#) are not physically able to perform either moderate or vigorous intensity exercise. Researchers created a study to learn more about how much exercise older adults are able to perform, and how that exercise affects their health.

The research team studied 6,489 female [participants](#) aged 63 to 99 years old. The researchers published their findings in the *Journal of the American Geriatrics Society*.

The participants agreed to take in-home exams, answer health questionnaires, and wear accelerometers (devices similar to fitness trackers). The participants also kept sleep logs.

The study was conducted between 2012 and 2013. The researchers reviewed death certificates as of September 2016 to learn how many participants had died.

At the beginning of the study, most participants were in their late 70s and most were considered overweight according to BMI standards (a ratio comparing height to weight). Nearly 30 percent were considered

obese.

Most participants scored 8.2 out of a possible 12 points on physical function assessments. Based on accelerometer measurements of the participants:

- 1 percent performed "low" light-intensity [physical activity](#)
- 29 percent performed "high" light-intensity physical activity
- 15 percent performed moderate to vigorous physical activity

After examining the deaths in the women according to their activity levels, the researchers learned that older women with higher levels of physical activity were less likely to die than women with lower levels of physical activity, no matter the cause of death.

The researchers concluded that their findings support encouraging [older women](#) to increase the amount of time they spend every day in light-intensity activity, and reduce the amount of time spent sitting.

More information: Michael J. LaMonte et al, Accelerometer-Measured Physical Activity and Mortality in Women Aged 63 to 99, *Journal of the American Geriatrics Society* (2017). [DOI: 10.1111/jgs.15201](#)

Provided by American Geriatrics Society

Citation: Light activity measured with fitness tracker linked to lower mortality in older women (2018, January 10) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-01-tracker-linked-mortality-older-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.