

Trastuzumab: No negative impact on cardiac function

January 3 2018

Long-term follow-up results of the NRG Oncology trial NSABP B-31 have shown that the addition of trastuzumab to adjuvant chemotherapy does not negatively affect cardiac function in women with node-positive, human epidermal growth factor receptor 2 (HER2)-positive, early-stage breast cancer who survive without cancer recurrence. Both patients who received trastuzumab in addition to chemotherapy and patients who received chemotherapy only maintained good cardiac function. Results of this follow-up analysis were published online in the *Journal of Clinical Oncology*.

"Breast cancer [patients](#) who received anthracycline and taxane-based [chemotherapy](#) with or without [trastuzumab](#) maintained excellent cardiac function a median of 8.8 years after treatment was started. In addition, patient reports of greater cardiac symptoms indicated more [cardiac problems](#) in the small group who had difficulties. Overall, for the relatively young [women](#) who entered this trial, the risks of cardiac late effects were minimal," commented Patricia A. Ganz, MD, of the University of California, Los Angeles, lead author of the article. Co-author Edward Romond of the University of Kentucky, Lexington, added, "Our study improved the 10 year survival for women with this type of aggressive [breast cancer](#) to 84% and now our long term follow-up of how these women are doing provides positive and encouraging data specifically about their cardiac function and health. We owe a great debt of thanks to the women who participated in this clinical trial."

Prior to this study, limited information was available regarding the long-

term follow-up of cardiac function and health-related quality of life in patients who remained disease-free following the completion of adjuvant chemotherapy. Previous studies suggested that early cardiac toxicity was a risk associated with adjuvant treatment when combined with trastuzumab. However, at median follow-up of almost 9 years, patients who were treated with anthracycline and taxane-based chemotherapy with added trastuzumab on NRG Oncology/NSABP B-31 did not experience long-term worsening of cardiac functioning or health-related quality of life.

Researchers assessed [cardiac function](#) through the measurement of left ventricular ejection fraction by multiple-gated acquisition (MUGA) scan and health-related quality of life using the Duke Activity Status Index (DASI) and the Medical Outcomes Study questionnaire. Current medications and comorbid conditions were also considered. Only 4.5% of patients from the control group and 3.4% from the trastuzumab group had a >10% decline in left ventricular ejection fraction from the baseline to a value

Citation: Trastuzumab: No negative impact on cardiac function (2018, January 3) retrieved 7 May 2024 from <https://medicalxpress.com/news/2018-01-trastuzumab-negative-impact-cardiac-function.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--