

Uneasiness in observers of unnatural android movements explained

January 9 2018



Fig.1. A movie of the android robot. Participants were shown movies of the android robot and those of the human model, and brain activity was compared between the two. Credit: Osaka University

It has been decades in the making, but humanoid technology has certainly made significant advancements toward creation of androids – robots with human-like features and capabilities. While androids hold

great promise for tangible benefits to the world, they may induce a mysterious and uneasy feeling in human observers. This phenomenon, called the "uncanny valley," increases when the android's appearance is almost humanlike but its movement is not fully natural or comparable to human movement. This has been a focus of study for many years; however, the neural mechanism underlying the detection of unnatural movements remains unclear.

The challenge prompted a team of Osaka University-centered researchers to delve deeper into the topic.

First, they recorded movies of Geminoid F, an [android](#) developed by Osaka University and Advanced Telecommunications Research Institute International, and its actual human model. This allowed the researchers to investigate what differences a slight degree of motion unnaturalness would make to feelings about the android.

Second, they used [functional magnetic resonance](#) imaging (MRI) to investigate what brain region activates when participants observed the android's and the human's movements in the movies, wherein they were only slightly different in naturalness. The [subthalamic nucleus](#) is found to be responsible for detecting the android's slight awkwardness.

"Visual observation of the android, compared with that of the human model, caused greater activation in the subthalamic nucleus (STN), which plays an important role in motor control," says Takashi Ikeda, lead author of the study, which was published recently in *Scientific Reports*.




Fig.2. Activation of the subthalamic nucleus. The movies of the android robot showed stronger activation in the subthalamic nucleus than those of the human model. The subthalamic nucleus is involved in the motor system and plays an important role in smooth motion. It is also a target area in deep brain stimulation to treat Parkinson disease. Credit: Osaka University

Notably, the STN is involved in the process of impairing dopaminergic neurons in the deep brain structure in Parkinson's disease patients. Deep brain stimulation to the STN dramatically improves rigidity or impairment of voluntary [movement](#) in these patients.

"Our data attest to commonalities between the movements of the android and Parkinson's disease patients. The android's movements were rigid and akinesic in a comparable way to the movements of a patient with mild Parkinson's disease," corresponding author Masayuki Hirata explains.

The current findings not only serve as valuable information for the design of androids, which could reduce the uncanny feeling in human observers, but also for elucidating the pathology of Parkinson's disease.

More information: Takashi Ikeda et al. Subthalamic nucleus detects unnatural android movement, *Scientific Reports* (2017). [DOI: 10.1038/s41598-017-17849-2](https://doi.org/10.1038/s41598-017-17849-2)

Provided by Osaka University

Citation: Uneasiness in observers of unnatural android movements explained (2018, January 9)
retrieved 7 September 2024 from <https://medicalxpress.com/news/2018-01-uneasiness-unnatural-android-movements.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.