

AIDS-defining events increase mortality risk, says study

February 9 2018, by Paul Govern

When they occur among people living with HIV, certain cancers and opportunistic infections are considered by health authorities as AIDS-defining events, or ADEs.

The Centers for Disease Control and Prevention recognizes 27 ADEs, from pneumonia to tuberculosis to cervical cancer to wasting syndrome. When a [death](#) is attributed to AIDS, usually one or more recognized ADEs is involved.

The availability of effective antiretroviral drug therapy has rendered HIV/AIDS a chronic disease in much of the world, no longer acutely fatal when treated. Among people living with HIV, at least in some developed regions, a so-called non-AIDS death appears at least as likely as a death attributed directly to AIDS.

In a study published recently in the *Journal of the International AIDS Society*, April Pettit, MD, MPH, assistant professor of Medicine, and colleagues focus on non-AIDS mortality among people on antiretroviral treatment in relatively high-income countries. They specifically examine mortality among patients who've survived an ADE since starting their HIV/AIDS [drug therapy](#).

The team examined records of 124,587 antiretroviral patients, with a median follow-up period of 5.2 years. Data were provided by the international Antiretroviral Therapy Cohort Collaboration (ART-CC).

The overall death rate was 9 percent. Among all deaths, 36 percent were coded as non-AIDS deaths, 24 percent were coded as AIDS deaths, and death codes were missing for the rest.

After adjusting their analysis for medical and demographic covariates (including time-updated covariates), the authors found that having survived an ADE is associated with 121 percent greater risk of non-AIDS death.

"These findings suggest that a common pathway may be independently driving both ADEs and NADE (non-AIDS) mortality. These findings underscore the need for future studies to elucidate a potential mechanism for this association, including that of chronic inflammation and immune activation due to ADEs," the authors wrote.

They focused on three events representing extremes of ADE severity: having survived tuberculosis, considered a mild ADE, is associated with 68 percent greater risk of non-AIDS death; having survived *Pneumocystis jirovecii* pneumonia (PJP), also considered mild, is associated with 121 percent greater risk of non-AIDS death; having survived non-Hodgkin lymphoma, considered severe, is associated with 195 percent greater risk of non-AIDS death.

Of these three ADEs, only TB was associated with a higher cardiovascular death rate. PJP was associated with higher rates of respiratory death and accident/suicide/overdose, with the latter association suggesting the influence of unmeasured confounders, according to the authors, such as mental illness and alcohol or drug abuse. Smoking and [body mass index](#), neither of which are measured by the ART-CC, are two other known confounders not included in the analysis.

The most frequent causes of non-AIDS death overall included cancer at

17 percent, accident/suicide/overdose at 16 percent, cardio-vascular disease also at 16 percent, non-AIDS infections at 15 percent, and liver disease at 13 percent.

Provided by Vanderbilt University

Citation: AIDS-defining events increase mortality risk, says study (2018, February 9) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-02-aids-defining-events-mortality-saysstudy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.