

Blood thinners may raise stroke risk in over-65s with kidney disease

February 14 2018



Micrograph showing cortical pseudolaminar necrosis, a finding seen in strokes on medical imaging and at autopsy. H&E-LFB stain. Credit: Nephron/Wikipedia

People over 65 years old may be increasing their stroke risk by taking anticoagulants for an irregular heartbeat if they also have chronic kidney disease, finds a new study led by UCL, St George's, University of London and the University of Surrey.

Based on their findings, published today in the *BMJ*, the researchers warn that doctors should be more cautious about prescribing anticoagulants, also known as [blood thinners](#), in this population until more studies can clarify the consequences of doing so.

"Chronic [kidney](#) disease is common among older people, and one in three people affected also have [atrial fibrillation](#), commonly called an [irregular heartbeat](#) - and for that, they typically get prescribed blood thinners to reduce their risk of stroke. We found that in this particular group, their medication seems to do the opposite of its intended effect," said the study's first author, Dr Shankar Kumar (UCL Centre for Medical Imaging).

The researchers estimate that close to half a million people over 65 in the UK have both [chronic kidney disease](#) and atrial fibrillation.

"People with chronic kidney disease tend to have numerous severe complications, including cardiovascular illnesses. As their blood clots more but they also bleed more easily, it is extremely difficult to strike a balance between different treatments," said senior author Professor John Camm, professor of clinical cardiology at St George's, University of London.

"As we found a paradoxical reduced mortality rate alongside increased rates of stroke and major bleeding, this is clearly a very complex area. We strongly call for randomised controlled studies to test the clinical value and safety of [anticoagulant](#) drug therapy for people with both atrial fibrillation and chronic kidney disease," said Dr Kumar.

In the population-based, [retrospective cohort study](#), the researchers used a Royal College of General Practitioners database to identify 4848 people over 65 with chronic kidney disease and newly diagnosed atrial fibrillation, half of whom were on anticoagulants and half were not. The

participants were monitored for a median of 506 days.

Over the study period, participants who were taking anticoagulants were 2.6 times as likely as those not on anticoagulants to have a stroke, and 2.4 times as likely to have a haemorrhage. The crude rates for ischaemic [stroke](#) (the most common type) were 4.6 per 100 person years for those on anticoagulants, and 1.5 for those not on blood thinners (in other words, if the participants were followed for exactly one year, there would be an estimated total of 4.6 ischaemic strokes per 100 people in the blood thinner group and 1.5 ischaemic strokes per 100 people not on [blood](#) thinners).

Mortality in the anticoagulant group was slightly lower; the researchers speculate it may have been due to a reduced risk of fatal strokes or heart attacks, but they say that more research is needed.

"Our work shows the power of big data in providing real world evidence to study important clinical scenarios. Until more data is available, general practitioners, nephrologists and cardiologists need to weigh up the risks and benefits of giving an anticoagulant and make a decision together with their patient," said co-author Professor David Goldsmith (Guy's and St Thomas' NHS Foundation Trust and St George's, University of London).

More information: *BMJ* (2018). [DOI: 10.1136/bmj.k342](https://doi.org/10.1136/bmj.k342)

Provided by University College London

Citation: Blood thinners may raise stroke risk in over-65s with kidney disease (2018, February 14) retrieved 2 May 2024 from <https://medicalxpress.com/news/2018-02-blood-thinners-over-65s-kidney-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.