


Learning about coronary heart disease from women

February 21 2018



Human heart. Credit: copyright American Heart Association

Coronary heart disease (CHD) is the number one cause of death in both men and women. While many risk factors of CHD, such as smoking, high blood pressure and age, are common among men and women, some metabolic risk factors, such as being diabetic, are more strongly

associated with increased risk of CHD in women than in men. Even though metabolic risk factors seem to be stronger in women, few studies have sought metabolic markers - that is, measurable indicators that can be detected in blood samples - of CHD in women. A new study by Brigham and Women's Hospital researchers recently addressed this gap, uncovering a new biomarker for CHD. Their results are published in *Circulation*.

"We were particularly interested in analyzing samples from women to understand some of the mechanisms underlying these risks," said corresponding author Nina P. Paynter, PhD, of the Department of Medicine at BWH. "We found a set of oxidized lipids that are novel markers for CHD, representing a different pathway contributing to CHD than the pathways we have previously considered."

By comparing the metabolic profiles of women with CHD and a control group of women, the researchers were able to identify one particular oxidized lipid as the strongest new risk marker for CHD. These results were replicated in a separate cohort of both men and women. Women in the highest quartile of the marker had a 4-fold higher risk of developing CHD in the next several years.

This marker is not associated with traditional risk markers, such as cholesterol levels and hypertension, suggesting that it contributes to CHD through a different pathway than the classical risk prediction markers. The exact mechanisms by which the identified oxidized lipid is contributing to CHD is unknown, and the researchers hope that future work will help to uncover how this risk marker is involved with CHD on a molecular level.

"Part of the reason that our study is so exciting is because it highlights that by focusing on specific patient groups, such as [women](#), that have been historically underrepresented in these sort of studies, we are able to

uncover new mechanisms may contribute to disease in the entire population," said senior author Kathryn M. Rexrode, MD, MPH, Chief of the Division of Women's Health of the Department of Medicine at BWH.

More information: Nina P. Paynter et al, Metabolic Predictors of Incident Coronary Heart Disease in Women, *Circulation* (2018). [DOI: 10.1161/CIRCULATIONAHA.117.029468](https://doi.org/10.1161/CIRCULATIONAHA.117.029468)

Provided by Brigham and Women's Hospital

Citation: Learning about coronary heart disease from women (2018, February 21) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-02-coronary-heart-disease-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.