

Differences in patient-reported pain between U.S. and European orthopedic patients

February 21 2018

American patients undergoing orthopedic surgical procedures report higher pain intensity scores compared to patients undergoing similar procedures in 13 European and non-European countries. American patients received more treatments for pain compared to their non-American counterparts, yet wished to have received even more treatment. However, most of the other assessments made by the American patients, e.g., patient-reported pain interference with function, were not different compared to those made by patients in other countries.

These findings are an outcome of analyzing over 14,000 patient questionnaires, collected within the world's largest acute pain registry, PAIN OUT, which is coordinated at the Jena University Hospital in Germany. "This survey is unique in that it compares data from a large number of patients from many different countries using highly standardized assessments," explains Ruth Zaslansky, first author of the study and researcher from the Department of Anesthesiology and Intensive Care Medicine at Jena University Hospital. Other authors included C. Richard Chapman, Professor emeritus from University of Utah in the US, and Winfried Meissner, coordinator of the PAIN OUT network from the Jena University hospital.

The authors say that to a certain extent, these results might be explained by the fact that in the U.S., more patients were on opioid medication prior to their surgery. This could reduce the therapeutic effect of analgesics after operations. However, also patients without preexisting

pain therapy report higher pain scores in the U.S. The authors then proposed two interpretations for their findings: Nurses in US hospitals typically ask patients far more often about the intensity of their pain compared to their colleagues in other countries. This might heighten awareness of pain, and interfere with distraction and other psychological coping mechanisms. If the treatment relies mostly on patients' pain intensity reports, this might lead to over-treatment. "The parameter 'pain intensity' might not be the most useful quality indicator for [pain management](#)," says Winfried Meissner.

Cultural differences regarding pain expectation

Another explanation could be cultural differences regarding pain expectation. It is possible that patients in the US are less willing to tolerate even [moderate pain](#) – an attitude reinforced by hospitals advertising the importance of patient satisfaction and by the promise of modern medicine to prevent any type of burden.

"The findings of this study suggest that 'worst pain' may possibly not be a good indicator for guiding pain therapy in individual [patients](#) or for evaluating quality of care in pain studies," emphasizes Ruth Zaslansky. The study findings do not support refraining from assessing pain but it is possible that other indicators, such as assessing the effect of [pain](#) on movement or on breathing, might be more useful in guiding treatment and mirroring quality of care. However, these hypotheses require further study.

More information: R. Zaslansky et al. Pain after orthopaedic surgery: differences in patient reported outcomes in the United States vs internationally. An observational study from the PAIN OUT dataset, *British Journal of Anaesthesia* (2018). [DOI: 10.1016/j.bja.2017.11.109](https://doi.org/10.1016/j.bja.2017.11.109)

Provided by Friedrich Schiller University of Jena

Citation: Differences in patient-reported pain between U.S. and European orthopedic patients (2018, February 21) retrieved 30 April 2024 from

<https://medicalxpress.com/news/2018-02-differences-patient-reported-pain-european-orthopedic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--