

# Exposure to chemicals used during fracking may cause pre-cancerous lesions in mice

February 7 2018, by Jeff Sossamon

---

Using more than 1,000 different chemicals, unconventional oil and gas (UOG) operations combine directional drilling and hydraulic fracturing, or "fracking," to release natural gas from underground rock. Today, researchers at the University of Missouri and the University of Massachusetts released a study that found that female mice exposed to mixtures of chemicals used in UOG operations during prenatal development had abnormal mammary glands in adulthood. Additionally, some of the mice developed pre-cancerous mammary lesions that may suggest they will be more sensitive to chemicals that cause cancer.

"Our earlier research showed that both male and [female mice](#) had alterations to hormone levels and reproductive organs resulting from exposures to these UOG contaminants," said Susan C. Nagel, an associate professor of Obstetrics, Gynecology and Women's Health in the School of Medicine. "We felt this could indicate that exposures to UOG chemical mixtures can produce a range of defects in animals exposed during vulnerable periods, such as development in the womb. So, we examined 23 UOG contaminants and compounds commonly used or produced in the fracking process."

In the study, female mice were exposed to various amounts of the 23 UOG chemicals from gestational day 11 to birth. Although no effects were observed on the mammary glands of these females prior to puberty, in early adulthood, female mice developed mammary lesions and hyperplasia, a condition that causes enlargement of an organ or tissue.

"We chose varying amounts of the UOG mixture in order to mimic a range of human exposures to these chemicals," Nagel said, who also serves as an adjunct associate professor of biological sciences in the MU College of Arts and Science. "These results suggest that the mammary [gland](#) is sensitive to mixtures of chemicals used in unconventional oil and gas production. Determining whether these fracking mixtures affect human populations is an important goal, particularly as the number of [fracking](#) sites within human population centers increases."

Additional, long-term studies are needed to evaluate these outcomes, the researchers said.

The study, "Prenatal [exposure](#) to unconventional oil and gas operation [chemical](#) mixtures altered mammary gland development in adult female [mice](#)" was published in the journal *Endocrinology*.

Provided by University of Missouri-Columbia

Citation: Exposure to chemicals used during fracking may cause pre-cancerous lesions in mice (2018, February 7) retrieved 13 May 2024 from <https://medicalxpress.com/news/2018-02-exposure-chemicals-fracking-pre-cancerous-lesions.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--