

# High detection rates of high-grade DCIS persist

February 7 2018

---



(HealthDay)—High detection rates of high-grade ductal carcinoma in

situ (DCIS) persist in consecutive subsequent screening rounds versus the prevalence round, while there are decreases in detection rates of low- and intermediate-grade DCIS, according to a study published in the February issue of *Radiology*.

Stefanie Weigel, M.D., Ph.D., from the University of Muenster in Germany, and colleagues compared grade-specific detection rates between the prevalence round (baseline screening) and two subsequent screening rounds of a population-based digital mammography screening program. There was a 22- to 30-month interval between all screening rounds.

The researchers found that cancer detection rates were lowest for low-grade DCIS over all screening rounds (range 0.11 to 0.25 per 1,000 women screened) and were highest for high-grade DCIS (range, 0.53 to 0.59 per 1,000 women screened). For low-grade DCIS, detection rates were significantly lower in the first and second screening rounds versus the prevalence round (odds ratios, 0.45 and 0.57, respectively; both P screening (odds ratios, 0.89 [P = 0.143] and 0.97 [P = 0.7], respectively).

"Grade-related changes of DCIS detection are suggestive of distinct dynamics of lesion progression," the authors write.

**More information:** [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: High detection rates of high-grade DCIS persist (2018, February 7) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-02-high-high-grade-dcis-persist.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.