

Study questions link between medical marijuana and fewer opioid deaths

February 6 2018

The association between medical marijuana and lower levels of opioid overdose deaths—identified previously in several studies—is more complex than previously described and appears to be changing as both medical marijuana laws and the opioid crisis evolve, according to a new RAND Corporation study.

The report—the most-detailed examination of [medical marijuana](#) and [opioid](#) deaths conducted to date—found that legalizing medical [marijuana](#) was associated with lower levels of opioid deaths only in states that had provisions for dispensaries that made medical marijuana easily available to patients. Opioid [death](#) rates were not lower in states that just provided legal protections to patients and caregivers, allowing them to grow their own marijuana.

In addition, the association between medical marijuana dispensaries and fewer opioid deaths appears to have declined sharply after 2010, when states began to tighten requirements on sales by dispensaries.

"Our findings are consistent with previous studies showing an association between the legalization of medical marijuana and lower deaths from overdoses of opioids," said Rosalie Liccardo Pacula, co-author of the study and co-director of the RAND Drug Policy Research Center.

"However, our findings show that the mechanism for this was loosely regulated medical marijuana dispensaries, and that the association between these laws and opioid mortality has declined over time as state

laws have more tightly regulated medical dispensaries and the opioid crisis shifted from prescription opioids to heroin and fentanyl," Pacula said. "This is a sign that medical marijuana, by itself, will not be the solution to the nation's opioid crisis today."

The study was published online by the *Journal of Health Economics*.

Researchers from RAND and the University of California, Irvine analyzed information about treatment admissions for addiction to pain medications from 1999 to 2012 and state-level overdose deaths from opioids from 1999 to 2013. They also identified state laws legalizing medical marijuana, examining provisions such as whether the regulations made marijuana easily accessible to patients by allowing dispensaries.

When the researchers narrowly focused on the time period from 1999 to 2010 and replicated a model used by other researchers, they obtained results similar to those previously published, showing an approximately 20 percent decline in [opioid overdose](#) deaths associated with the passage of any state medical marijuana law. However, these general findings were driven by states that had laws allowing for loosely regulated marijuana dispensary systems.

When researchers extended their analysis through 2013, they found that the association between having any medical marijuana law and lower rates of opioid deaths completely disappeared. Moreover, the association between states with medical marijuana dispensaries and opioid mortality fell substantially as well.

The researchers provide two explanations for the decline in the association between medical marijuana dispensaries and opioid harm. First, states that more recently adopted laws with medical marijuana dispensaries more tightly regulated them, in response to a U.S. Justice Department memo saying it would not challenge state-level [medical](#)

[marijuana laws](#) so long as dispensary sales were in full compliance with state regulations. Second, beginning in 2010, the primary driver of the opioid crisis and related deaths became illicit opioids, mainly heroin and then fentanyl, not prescription opioids.

The study also found no evidence that states with medical marijuana laws experience reductions in the volume of legally distributed opioid analgesics used to treat pain. Even if medical marijuana patients were substituting medical marijuana for opioids in medical marijuana states, these patients did not represent a measurable part of the medical opioid analgesic market.

"While our study finds that medical marijuana dispensaries reduce some of the harms associated with the misuse of opioids, there is little evidence that this is happening because a large number of patients suffering from pain are using marijuana instead of opioid medications," Pacula said. "Either the [patients](#) are continuing to use their opioid pain medications in addition to marijuana, or this patient group represents a small share of the overall medical opioid using population."

The RAND study was conducted before any any states had begun to allow retail sales of recreational marijuana.

"Our research suggests that the overall story between medical marijuana and opioid deaths is complicated," Pacula said. "Before we embrace marijuana as a strategy to combat the opioid epidemic, we need to fully understand the mechanism through which these laws may be helping and see if that mechanism still matters in today's changing opioid crisis."

Provided by RAND Corporation

Citation: Study questions link between medical marijuana and fewer opioid deaths (2018,

February 6) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-02-link-medical-marijuana-opioid-deaths.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.