

Study investigates the molecular basis for male predisposition to autism

February 7 2018

Exposure to androgens (male hormones) during brain development alters genes related to autism spectrum disorder (ASD), according to a new study published in *Biological Psychiatry*. Using male human cells, researchers at the University of Strasbourg, France, identified key genes that are regulated by testosterone and that contribute to the risk for autism, generating important insight into how male hormones might contribute to the increased male susceptibility to ASD.

The findings provide a clue as to why ASD affects boys about four times as much as girls. Male fetuses produce androgens during critical stages of brain development when cells divide and develop into neurons. The researchers showed that androgens increase the spread of cells and prevent them from death, which could predispose boys to ASD by contributing to the excessive brain growth that occurs in people with ASD during the first years of life.

"Understanding the mechanisms for the male preponderance for autism is like pulling on a loose thread. It could help to 'unravel' important mechanisms contributing to autism risk," said John Krystal, Editor of *Biological Psychiatry*. Identifying a role of androgens in this risk could be important for prevention or the development of potential treatments for ASD.

In the study, co-senior authors Jean-Louis Mandel, M.D., Ph.D., and Amélie Piton, Ph.D., and colleagues used human pluripotent neural stem cells to model the [cells](#) that generate neurons during [brain](#)

[development](#). Treatment with the testosterone metabolite DHT led to subtle changes in the expression of about 200 [genes](#), several of which have previously been associated with ASD. Some of the genes that were most affected by DHT treatment included NRCAM, which has been linked to the brain abnormalities and symptoms in ASD, and FAM107A, which is increased in people with ASD. FAM107A also appeared to play a part in androgen's ability to increase cell numbers in the study.

"These effects of [male hormones](#) may therefore contribute to the increased sensitivity of the male [brain](#) to develop ASD when also exposed to other genetic or environmental factors," said Dr. Piton, suggesting that the biological explanation for the sex-specific inequality of ASD points to a predisposition in males, rather than a protective effect in females.

In addition to providing a clue into the male susceptibility to ASD, Dr. Piton says that the list of genes that were altered by androgens in the study might be useful to identify new genes that might be involved in ASD or in other diseases that occur more often in males.

More information: Angélique Quartier et al. Genes and pathways regulated by androgens in human neural cells, potential candidates for the male excess in autism spectrum disorders, *Biological Psychiatry* (2018). [DOI: 10.1016/j.biopsych.2018.01.002](https://doi.org/10.1016/j.biopsych.2018.01.002)

Provided by Elsevier

Citation: Study investigates the molecular basis for male predisposition to autism (2018, February 7) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-02-molecular-basis-male-predisposition-autism.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.