

Are older adults with knee pain less active than the general population?

February 22 2018



Credit: CC0 Public Domain

A new *Arthritis Care & Research* study found that moderate-to-vigorous physical activity levels are similarly low in older adults with symptomatic knee osteoarthritis and those from the general population without osteoarthritis or knee pain.

Because the general population is doing as little as individuals with [knee pain](#) that may hinder activity, the findings point to the need for efforts to increase [physical activity](#) for all.

"We were a little surprised to see similar low levels of physical activity in both those with and without painful knee arthritis. I think this is a wake-up call to everyone that we all need to be doing more activity," said senior author Dr. Daniel White, of the University of Delaware.

More information: Louise M. Thoma et al, Are older adults with symptomatic knee osteoarthritis less active than the general population?: Analysis from the Osteoarthritis Initiative and NHANES, *Arthritis Care & Research* (2018). [DOI: 10.1002/acr.23511](https://doi.org/10.1002/acr.23511)

Provided by Wiley

Citation: Are older adults with knee pain less active than the general population? (2018, February 22) retrieved 10 April 2024 from <https://medicalxpress.com/news/2018-02-older-adults-knee-pain-population.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
