

Patients with kidney disease with heart defibrillators at greater risk of hospitalization

February 5 2018

In a study of nearly 6,000 community-based patients with chronic kidney disease and heart failure, the use of implantable cardioverter defibrillators was associated with a significantly increased risk of subsequent hospitalization.

The study, "Long Term Outcomes Associated with Implantable Cardioverter Defibrillator in Adults with Chronic Kidney Disease," was conducted by researchers from Kaiser Permanente and the University of Washington. It was published today in *JAMA Internal Medicine*.

"The finding surprised us," said lead author Nisha Bansal, MD, MAS, of the Kidney Research Institute of University of Washington's Division of Nephrology. "Chronic [kidney](#) disease is common in adults with heart failure, and is associated with a greater risk of heart attack. However, in this observational study we did not find a significant overall benefit from ICDs for [patients](#) with kidney disease."

ICDs are small implantable devices that deliver electrical shocks to the heart in order to help control life-threatening irregular heartbeats, or arrhythmias. (Pacemakers are a different implanted device, that gives off low-energy electrical pulses to treat less-dangerous heart rhythms.) ICDs are commonly used to reduce the likelihood of sudden death in patients at high risk of cardiac arrest.

"Because placing an ICD can be expensive and can cause other complications, it is critical to better understand how to best use this therapy in high-risk patients, such as those with chronic kidney disease," said senior author Alan S. Go, MD, a research scientist with the Kaiser Permanente Northern California Division of Research. "Based on our results, clinicians should carefully consider the risks and benefits of ICDs when recommending them for this subset of patients."

Chronic kidney disease affects an estimated 14 percent of adults in the United States, and heart failure is the leading cause of death among them. About 30 percent of the estimated 5.7 million U.S. adults with heart failure also have chronic kidney disease, explained Dr. Bansal. The researchers identified 5,877 adults with chronic kidney disease and heart failure with reduced left ventricular ejection fraction, all Kaiser Permanente patients in California, Northwest and Colorado regions, between 2005 and 2013; among them, 1,556 had an ICD, and 4,321 did not have an ICD.

After accounting for differences in demographics, numerous other health conditions, the use of heart medications, and confounding risk factors, there was no difference in mortality between patients with chronic kidney disease who did or did not receive an ICD. However, ICD placement was associated with a 49 percent increased risk of subsequent hospitalization due to [heart](#) failure and a 25 percent increased risk of hospitalization for any reason.

Clinical trials have examined the value of ICDs in the general [heart-failure](#) population, Dr. Go noted, and have shown that the device increases patients' protection against sudden cardiac death and improves their overall survival. "The data from our study certainly need to be confirmed by clinical trials conducted specifically in patients with [chronic kidney disease](#)," he added.

Provided by Kaiser Permanente

Citation: Patients with kidney disease with heart defibrillators at greater risk of hospitalization (2018, February 5) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-02-patients-kidney-disease-heart-defibrillators.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.