


Poor fitness linked to weaker brain fiber, higher dementia risk

February 14 2018, by James Beltran



Brain imaging shows yellow and reddish pixels representing areas where the

functionality of white matter is associated with higher fitness levels. The images are based on cumulative data from patients in a study showing potential links between physical fitness and deterioration of white matter. Credit: UT Southwestern Medical Center

Scientists have more evidence that exercise improves brain health and could be a lifesaving ingredient that prevents Alzheimer's disease.

In particular, a new study from UT Southwestern's O'Donnell Brain Institute suggests that the lower the [fitness level](#), the faster the deterioration of vital nerve fibers in the brain. This deterioration results in cognitive decline, including memory issues characteristic of dementia patients.

"This research supports the hypothesis that improving people's [fitness](#) may improve their brain health and slow down the aging process," said Dr. Kan Ding, a neurologist from the Peter O'Donnell Jr. Brain Institute who authored the study.

White matter

The study published in the *Journal of Alzheimer's Disease* focused on a type of brain tissue called white matter, which is comprised of millions of bundles of [nerve fibers](#) used by neurons to communicate across the brain.

Dr. Ding's team enrolled older patients at high risk to develop Alzheimer's disease who have early signs of memory loss, or mild cognitive impairment (MCI). The researchers determined that lower fitness levels were associated with weaker white matter, which in turn correlated with lower [brain function](#).

Distinctive tactics

Unlike previous studies that relied on study participants to assess their own fitness, the new research objectively measured cardiorespiratory fitness with a scientific formula called maximal oxygen uptake. Scientists also used brain imaging to measure the functionality of each patient's [white matter](#).

Patients were then given memory and other cognitive tests to measure brain function, allowing scientists to establish strong correlations between [exercise](#), [brain health](#), and cognition.

Lingering mysteries

The study adds to a growing body of evidence pointing to a simple yet crucial mandate for human health: Exercise regularly.

However, the study leaves plenty of unanswered questions about how fitness and Alzheimer's disease are intertwined. For instance, what fitness level is needed to notably reduce the risk of dementia? Is it too late to intervene when patients begin showing symptoms?

Some of these topics are already being researched through a five-year national clinical trial led by the O'Donnell Brain Institute.

The trial, which includes six medical centers across the country, aims to determine whether regular aerobic exercise and taking specific medications to reduce high blood pressure and cholesterol levels can help preserve brain function. It involves more than 600 older adults at high risk to develop Alzheimer's disease.

"Evidence suggests that what is bad for your heart is bad for your brain.

We need studies like this to find out how the two are intertwined and hopefully find the right formula to help prevent Alzheimer's disease," said Dr. Rong Zhang of UT Southwestern, who oversees the clinical trial and is Director of the Cerebrovascular Laboratory in the Institute for Exercise and Environmental Medicine at Texas Health Presbyterian Hospital Dallas, where the Dallas arm of the study is being carried out.

Prior findings

The research builds upon prior investigations linking healthy lifestyles to better brain function, including a 2013 study from Dr. Zhang's team that found neuronal messages are more efficiently relayed in the brains of older adults who exercise.

In addition, other teams at the O'Donnell Brain Institute are designing tests for the early detection of patients who will develop dementia, and seeking methods to slow or stop the spread of toxic proteins associated with the [disease](#) such as beta-amyloid and tau, which are blamed for destroying certain groups of neurons in the [brain](#).

"A lot of work remains to better understand and treat dementia," said Dr. Ding, Assistant Professor of Neurology & Neurotherapeutics. "But, eventually, the hope is that our studies will convince people to exercise more."

More information: Kan Ding et al. Cardiorespiratory Fitness and White Matter Neuronal Fiber Integrity in Mild Cognitive Impairment, *Journal of Alzheimer's Disease* (2017). [DOI: 10.3233/JAD-170415](https://doi.org/10.3233/JAD-170415)

Provided by UT Southwestern Medical Center

Citation: Poor fitness linked to weaker brain fiber, higher dementia risk (2018, February 14)
retrieved 2 May 2024 from
<https://medicalxpress.com/news/2018-02-poor-linked-weaker-brain-fiber.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.