

New study shows repurposing leukemia drugs may prevent melanoma metastasis

February 26 2018, by Allison Perry



A new study from Markey researchers shows that repurposing drugs used to treat leukemia has promise for preventing melanoma metastasis. Credit: University of Kentucky

Data from a new study led by University of Kentucky Markey Cancer Center researchers shows that repurposing drugs used to treat leukemia



has promise for preventing melanoma metastasis.

Published in *Science Signaling*, the study showed new evidence linking the activation of ABL kinases – cancer-promoting genes – to the secretion of pro-metastatic cathepsins in <u>melanoma</u>. Cathepsins are enzymes that degrade proteins and are highly expressed in cancer cells, resulting in their release into the environment between the cells. These enzymes "chew up" the fibrous matrix around tumors, which allows them to get into the blood stream and lymphatic system and spread around the body.

Their work showed that ABL kinases induce cathepsin expression and secretion by increasing the activity of key <u>transcription factors</u> that upregulate numerous proteins involved in metastasis. Transcription factors bind to the regulatory part of genes and induce their expression. This study is the first to demonstrate that ABL kinases not only increase the abundance of the transcription factors, but also regulate the ability of these transcription factors to bind to the promoters and induce gene expression.

Lastly, the researchers found that ABL kinases inhibitors already approved by the Food & Drug Administration (FDA) for treating leukemia also prevented metastasis induced by secreted cathepsins in animal models of metastatic melanoma.

"These data have important therapeutic implications, as drugs that inhibit ABL kinases have been used for decades to treat leukemia with few side effects," said Rakshamani Tripathi, postdoctoral scholar in the UK Department of Pharmacology and Nutritional Sciences and lead author on the study. "Repurposing these drugs may represent a new strategy for targeting cathepsins and may be an effective approach for treating metastatic melanoma patients."



The number of new cases of melanoma, as well as melanoma death rates, have increased steadily for the past 30 years in the United States. Metastasis to distant organ sites such as the lung, liver brain and bone is the major cause of melanoma-related deaths. Despite new advances in metastatic melanoma treatment, the 5-year survival rate is only 15 to 20 percent.

Provided by University of Kentucky

Citation: New study shows repurposing leukemia drugs may prevent melanoma metastasis (2018, February 26) retrieved 27 April 2024 from <u>https://medicalxpress.com/news/2018-02-repurposing-leukemia-drugs-melanoma-metastasis.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.