


'Social brain' networks are altered at a young age in autism

February 27 2018



Each dot represents the gaze position for an individual child watching the movie. The blue dots on the left represent the typically developing toddlers. The red dots represent toddlers with Autism Spectrum Disorders (ASD). The face wasn't blurred during the experiment. Credit: UNIGE

As infants develop, they preferentially move towards and respond to social cues - such as voices, faces and human gestures. At the same time, their brain develops a network of regions that specialise in translating these cues, known as the 'social brain'. However, a common observation in infants later diagnosed with Autism Spectrum Disorders (ASD) is reduced sensitivity and attention towards these social cues during the first year of life. This apparent indifference to social cues is thought to

ultimately hinder the normal development of the social brain at early developmental stages. A team of researchers from the University of Geneva (UNIGE), Switzerland, now brings direct evidence of this hindered development during the toddler to pre-school years in autism. Their findings are published in the journal *eLife*.

The results of their study suggest that interventions targeting children's ability to respond to [social cues](#) at this critical early age could rewire the [brain](#) while it is still possible to do so, potentially restoring social brain development. Dr Holger Sperdin, Postdoctoral Research Associate at the UNIGE's Faculty of Medicine and lead author of the study, explains what he and his team set out to discover: "As toddlers with ASD have less preferential attention for social cues, we hypothesised that when we showed them moving social images, they would demonstrate differences in both the way they visually explore these images and in the way their brain networks process social information, compared with typically developing toddlers."

Eye-tracking technology

The team used an electrophysiological monitoring method called electroencephalography (EEG) to study the children's brain activity, and powerful [eye-tracking technology](#) to observe their gaze while they watched movies featuring human social interactions. They found that the children with ASD had different gaze patterns while watching the movies to the typically developing infants, and that this was accompanied by alterations in nerve cell connectivity and information flow in the brain.

In those with ASD, the team also observed what is known as 'increased driving' in two specific frequencies of brain waves - alpha and theta - as well as high levels of connectivity between nerve cells in certain regions in the brain. The theta brain wave frequency and the regions of the brain

affected are both known to be important components of the 'social brain', and the alpha frequency is important for visual attention.

These findings represent the first evidence that differences in the visual exploration of images coincide with changes in connectivity between key regions of the social brain in very young children with ASD. Brain regions generating these brain wave frequencies may therefore develop differently in children with ASD compared with their typically developing peers.

"Our results show for the first time the presence of alterations in information flow from brain areas involved in social cue processing in toddlers and pre-schoolers with Autism Spectrum Disorders (ASD)," concludes senior author Professor Marie Schaer, Assistant Professor at the University of Geneva, Switzerland. "These alterations within regions of the social brain are present at early stages of ASD and justify further investigation into whether therapeutic interventions targeting social orienting skills may help to remediate social brain development during this critical stage when neural plasticity is still possible."

More information: Holger Franz Sperdin et al, Early alterations of social brain networks in young children with autism, *eLife* (2018). [DOI: 10.7554/eLife.31670](https://doi.org/10.7554/eLife.31670)

Provided by University of Geneva

Citation: 'Social brain' networks are altered at a young age in autism (2018, February 27) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-02-social-brain-networks-young-age.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.