

## Spinal fusion surgery for lower back pain is costly, unsupported by evidence

February 20 2018, by Gustavo Machado, Christine Lin And Ian Harris



Credit: Karolina Grabowska from Pexels

Back pain affects <u>one in four</u> Australians. It's so common, nearly all of us (<u>about 85%</u>) will have at least one episode at some stage of our lives. It's one of the most common reasons <u>to visit a GP</u> and the main health



condition forcing older Australians to <u>retire prematurely</u> from the workforce.

Treatment costs for back pain in Australia total almost <u>A\$5 billion</u> every year. A great proportion of this is spent on spinal surgical procedures. Recently, Choosing Wisely, the campaign to educate medical professionals and the public about tests, treatments and procedures that have little benefit, or lead to harm, <u>added spinal fusion</u> for lower back pain to its list.

This is because, despite rates of the procedure being on the rise, current evidence doesn't support spinal <u>fusion</u> for back pain. Randomised trials (regarded as studies providing the highest-quality evidence) suggest <u>spinal fusion has little advantage</u> over a well-structured rehabilitation program, or psychological interventions, for back pain.

## What is spinal fusion?

Spinal surgery is most commonly performed to remove pressure on nerves that causes pain and other nerve symptoms in the legs. This surgery is called decompression. The next most common procedure is spinal fusion, where two or more vertebrae are joined together (using such methods as transplanted bone from the patient, a donor or artificial bone substitutes) to stop them moving on each other and make one solid bone.

Spinal fusion may be performed for fractures, dislocations and tumours, and is commonly performed in conjunction with decompression. For back pain, it's performed when the origin of the pain is thought to be related to abnormal or painful movement between the vertebrae (from degenerative joints and discs, for example).

Rates of spinal fusions have been rising and continue to increase,



outstripping other surgical procedures for back pain. In the United States, rates of spinal fusion more than doubled from 2000 to 2009. In Australia, rates <u>increased</u> by 167% in the <u>private sector</u> between 1997 and 2006, despite almost no increase in the public sector.

Spinal fusion rates <u>differ significantly</u> between regions of Australia, with the highest being in Tasmania and the lowest in South Australia: a sevenfold variation. Significant variations are also seen between countries. For instance, spinal fusion rates in the United States are <u>eight times</u> those in the United Kingdom.

The greatest increase in the use of spinal fusion has been in <u>older</u> <u>Australians</u>, often in conjunction with decompression surgery for <u>spinal</u> <u>stenosis</u> – a condition that causes narrowing of the <u>spinal canal</u> (the cavity that runs through the <u>spinal cord</u>).

Differences in clinical training, professional opinion, and local practices are likely to play a role in such variations.

## **Evidence for spinal surgery**

There is little high-quality evidence to support the use of spinal fusion for most back-related conditions, including <u>spinal stenosis</u>. And there is <u>disagreement between surgeons</u> on when <u>spinal fusion surgery</u> should be performed, not only for back <u>pain</u> but also for more acute conditions such as tumours and spine fractures.

There have also been no studies comparing spinal fusion to a placebo procedure. Most research to date compares one fusion technique to <u>another technique</u> or to a form of <u>non-surgical treatment</u>, so we still don't know whether <u>spine fusion</u> is effective against placebo.

We also know that spine fusion surgery is expensive and associated with



<u>more complications</u> than decompression surgery. And the surgery often fails. Around <u>one in five</u> patients who undergo spine fusion will have revision surgery within ten years.

<u>Research also shows</u> most patients having spine fusion surgery under workers' compensation won't return to the usual job, will still be having physiotherapy and be on opioid medication two years after surgery.

## So why are rates going up?

There are <u>several factors</u>, including an ageing population, that may contribute to the rapid increase in spinal fusion despite the lack of evidence supporting its use. Financial incentives might also explain the differences in rates between private and public sectors in Australia and between the United Kingdom and the United States.

We don't have high-quality evidence on the benefits and harms of <u>spinal</u> <u>fusion</u>. This means there is uncertainty, which allows practitioners to continue doing the procedures they were trained to do unchallenged. This then leads to overtreatment, particularly where reimbursement rates are high, such as in the <u>workers' compensation</u> setting.

Uncertainty about the appropriateness of spine fusion results in practice variation, wastes scarce health care resources and leads to worse patient outcomes.

We need better research in this area. This means research efforts should shift from studies looking at different ways of performing the <u>surgery</u> and focus on investigating whether or not it works better than nonoperative treatments or a placebo, and, if so, whether the benefits outweigh the harms.

In the absence of such evidence, patients can consider other evidence-



based and less costly treatments, such as <u>exercise</u>, <u>cognitive behavioural</u> <u>therapy and physiotherapy</u>.

This article was originally published on <u>The Conversation</u>. Read the <u>original article</u>.

Provided by The Conversation

Citation: Spinal fusion surgery for lower back pain is costly, unsupported by evidence (2018, February 20) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2018-02-spinal-fusion-surgery-pain-costly.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.