

Younger age at diabetes diagnosis is linked to higher risk of death from heart disease and stroke

February 22 2018

While type 2 diabetes (T2D) was once considered a disease largely confined to older people, the global epidemic of obesity and overweight has seen diagnoses rocket in young adults, adolescents and even appear in young children. New research published in *Diabetologia* (the journal of the European Association for the Study of Diabetes [EASD]) shows that the earlier a person is diagnosed with T2D, the higher their risk of death from heart disease and stroke, but, unusually, the lower their risk of death from cancer.

In almost all countries of the world, diabetes rates are increasing substantially in younger adults, aged 20-45 years. Rates are also continuing to increase in adults over 45 years old, however not as sharply as in younger adults. The increase in the [younger adults](#) means there is a steadily growing pool of [diabetes patients](#) who are exposed to diabetes for a longer period in their lives.

The study by Professor Dianna Magliano and Professor Jonathan Shaw (Baker Heart and Diabetes Institute, Melbourne, Australia) and colleagues analysed the data of 743,709 Australians with T2D who were registered on Australia's National Diabetes Services Scheme (NDSS) over a 15-year period between 1997 and 2011. All-cause [mortality](#) and mortality due to cardiovascular disease (CVD), cancer and all other causes were identified.

The average (median) age at T2D [diagnosis](#) was 59 years, and a total of 115,363 deaths occurred during the study period. The authors say: "An earlier diagnosis of type 2 diabetes—and thus a longer duration of disease—was associated with a higher risk of all-cause mortality, primarily driven by [cardiovascular disease](#) (CVD) mortality."

The data showed that for two people of the same age, the one with a 10-year earlier diagnosis (equivalent to 10 years' longer duration of diabetes) had a 20% to 30% increased risk of all-cause mortality and about a 60% increased risk of CVD mortality. The effects were similar in men and women.

authors say: "Evidence is accumulating to suggest that earlier onset of type 2 diabetes is associated with an increased risk of complications and comorbidities compared with later onset, and that the development and progression of complications might be more aggressive in those with earlier onset."

They add: "As such, increased clinical attention is imperative for individuals with earlier-onset type 2 diabetes. Efforts should focus on timely optimisation of individuals' self-management skills and medical treatment to prevent or reduce the onset of complications and comorbidities. Additionally, there is a need to identify and screen those at high risk of developing diabetes so that individuals can make lifestyle changes that will prevent or delay the onset of diabetes."

Other interesting findings from the study by Professors Magliano, Shaw and colleagues include that for mortality due to cancer (all cancers and colorectal and lung cancers), earlier diagnosis of type 2 diabetes was associated with lower mortality compared with diagnosis at an older age. While this may appear unusual, the authors point out that "it is possible that following a diagnosis of [diabetes](#), people have more frequent contact with the healthcare system, which may increase the likelihood of any

present but undiagnosed [cancer](#) being detected."

More information: *Diabetologia* (2018). link.springer.com/article/10.1007/s00125-018-4544-z

Provided by Diabetologia

Citation: Younger age at diabetes diagnosis is linked to higher risk of death from heart disease and stroke (2018, February 22) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-02-younger-age-diabetes-diagnosis-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.