

Cardiovascular health disparities between whites and minorities narrow, study shows

March 21 2018, by Enrique Rivero



Credit: Wikimedia Commons

The nation's overall cardiovascular health worsened from 1988 to 2014, with disparities among racial and ethnic groups dropping slightly. But the reduction in disparities was due to worsening health among whites—not

improvements among African-Americans and Mexican-Americans, a new UCLA-led study suggests.

"The reason for the reduction in disparities was unexpected," said lead author Dr. Arleen Brown, professor of medicine in the division of general internal medicine at the David Geffen School of Medicine at UCLA and Olive View Medical Center. "Whites were the only group we studied where the prevalence of optimal cardiovascular [health](#) declined consistently over the time period."

Among the findings:

- From 1988 to 2014, the percentage of whites with optimal cardiovascular health fell 15.3 percentage points among those ages 25 to 44, and 4.6 percentage points among those 65 and older.
- By contrast, the percentage of African-Americans with optimal cardiovascular health fell 3 percentage points among those ages 25 to 44, and 0.3 percentage points among those 65 and older.
- The percentage of U.S.-born Mexican-Americans with optimal health fell 8.6 percentage points among those ages 25 to 44, and 2.6 percentage points for those over 65. For foreign-born people of Mexican descent, the percentage-point drops were 7.6 and 1.3, respectively.

The researchers used National Health and Nutrition Examination Survey data from 1988 through 2014 to examine trends—by race, ethnicity and country of birth—of the seven controllable factors affecting heart health: blood pressure, cholesterol, blood glucose, weight, physical activity, diet and smoking. They then broke down the data into five time periods to more easily see trend changes and to draw a sharper contrast between the beginning and end of the total time span.

The findings are published in the peer-reviewed *Annals of Internal Medicine*.

Brown acknowledged that the study may be limited by not being representative of the total U.S. population, possibly because of under-reporting of heart disease, a lack of subgroups among the racial and [ethnic groups](#), and other factors. But they are a start toward a more complete picture of [cardiovascular disease](#) in the United States, and based on the data analysis, could provide direction for future interventions, the study stated.

"Cardiovascular disease disproportionately affects minorities at an earlier age than whites, and contributes to the economic burden of [health disparities](#)," Brown said. "There are ways to modify both biological and [behavioral risk factors](#) that lead to [heart disease](#). This data can help prioritize those intervention strategies."

More information: Arleen F. Brown et al. Trends in Racial/Ethnic and Nativity Disparities in Cardiovascular Health Among Adults Without Prevalent Cardiovascular Disease in the United States, 1988 to 2014, *Annals of Internal Medicine* (2018). [DOI: 10.7326/M17-0996](https://doi.org/10.7326/M17-0996)

Provided by University of California, Los Angeles

Citation: Cardiovascular health disparities between whites and minorities narrow, study shows (2018, March 21) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-03-cardiovascular-health-disparities-whites-minorities.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.