

Climate change promotes the spread of mosquito and tick-borne viruses

March 16 2018

Spurred on by climate change, international travel and international trade, disease-bearing insects are spreading to ever-wider parts of the world.

This means that more humans are exposed to viral infections such as Dengue fever, Chikungunya, Zika, West Nile fever, Yellow fever and Tick-borne encephalitis.

For many of these diseases, there are as yet no specific antiviral agents or vaccines.

Global warming has allowed [mosquitoes](#), ticks and other disease-bearing insects to proliferate, adapt to different seasons, migrate and spread to new niche areas that have become warmer.

These are the findings of a JRC report that aims to raise awareness about the threat posed by the spread of arboviruses (arthropod-borne viruses).

The growing spread of arboviruses

Aedes mosquitoes spread several arboviruses, including Dengue, Chikungunya, Zika, West Nile and Yellow fever viruses.

These mosquitoes thrive in urban settings due to the lack of natural predators and the ready availability of food and habitats in which to

procreate.

They have existed in Africa and Asia for many years and are now becoming more and more widespread.

They have recently become established in some European countries and the Americas, largely as a result of international travel and trade.

Their alarming spread poses a problem for public health. They are difficult to eradicate - their larvae can survive for months, even in suboptimal humidity and temperature conditions.

The tick-borne encephalitis virus (TBEV) has been found in several European countries, including Austria, the Czech Republic, Germany, Lithuania, Poland, Slovakia, Sweden and, more recently, the Netherlands.

One of the more recently reported vectors for the virus, the *Dermacentor reticulatus* tick species, is rapidly spreading through Europe. It has a high reproduction rate, is cold resistant and can live underwater for months.

Humans can be infected by a tick bite or through consuming unpasteurised dairy products that do not meet EU safety standards and have come from infected animals. Luckily, TBEV can be vaccinated against.

Zika virus - a serious concern for Europe

Zika virus has received a lot of media attention due to its association with neurological disorders such as Guillain-Barré Syndrome (GBS) and the development of microcephaly (abnormally small head) in foetuses.

It is difficult to diagnose and there is no cure or vaccine. First identified

in 1947 in the Zika forest of Uganda, its spread is a serious concern given the growing presence of its main vector, the mosquito *Aedes albopictus*, in temperate zones including Europe and America.

The first documented outbreak of ZIKV infection was reported in 2007 in Micronesia. Since then it spread to French Polynesia and Brazil, where it infected up to 1.3 million people in 2015.

More than 70 territories worldwide have confirmed autochthonous (indigenous) cases of ZIKV. By March 2017, 2 130 Europeans were reported to have travel-associated ZIKV infections.

Mosquito control strategies

The report describes and discusses several methods that have been used to control the spread of mosquitoes, including insecticides, mosquito traps, genetic modification, land reclamation and habitat surveillance.

Currently, the safest and most readily available and effective methods of controlling mosquitoes are mosquito traps (for relatively small areas) and nets, and the reduction of potential breeding sites (standing water).

While the research team behind the study advocate better control of mosquito populations, they also warn that it would be unwise to remove mosquitoes completely from the ecosystem.

They are part of the food web for some species, and pollinate many plants. Wiping them out completely could have negative effects on nature, and consequently on humans.

Provided by European Commission Joint Research Centre

Citation: Climate change promotes the spread of mosquito and tick-borne viruses (2018, March 16) retrieved 2 May 2024 from <https://medicalxpress.com/news/2018-03-climate-mosquito-tick-borne-viruses.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.