

New colon cancer finding could lead to earlier diagnosis—and better outcomes

March 28 2018



Cancer—Histopathologic image of colonic carcinoid. Credit: Wikipedia/CC BY-SA 3.0

For many years, physicians have puzzled over why people with "clean" colonoscopies went on to develop colon cancer. New findings from the Oklahoma Medical Research may help explain why, and the discovery could lead to ways to detect these cancers earlier and more effectively.

Trailing only [lung cancer](#), [colon cancer](#) is the leading cause of [cancer](#) deaths for men and women, killing 65,000 Americans each year. Still, life expectancy improves considerably if the cancer is detected early: People whose colon cancer is discovered in the earliest stage have a 5-year survival rate of 90 percent, while those whose cancer is found in the latest stage have an 8 percent rate.

The most common method of screening is a colonoscopy, where doctors use a flexible scope to examine the colon. However, certain cancer-causing [polyps](#) can be easily missed during these examinations.

"Some polyps are embedded in the surface of the colon, and they're also flat and covered up," said David Jones, Ph.D., who holds the Jeanine Rainbolt Chair for Cancer Research at OMRF. "This makes them incredibly difficult for doctors to detect."

For a long time, said Jones, it was thought that colon cancers that developed in patients who had clean colonoscopies were coming about through some unknown mechanism that didn't involve polyps. "Now it is clear that these hidden polyps might be responsible for up to 30 to 40 percent of colon cancers that develop later," he said.

Working with a team of researchers, Jones analyzed the genetic composition of the hidden polyps. "Most cancers—and most polyps—need more than one mutation to form. However, in these polyps, only one gene, called BRAF, was mutated," he said.

Because of these telltale markers identifying the polyps, Jones said, it would be possible to create a diagnostic test to analyze fecal samples to look for these changes prior to a colonoscopy. "If changes are present, it would be a way for the doctors to know to look for a hiding polyp," said Jones.

Further analysis by the researchers also showed that the mutation caused a wave of alterations in the DNA. "It's probably the change in BRAF combined with these changes that leads to polyps forming."

Jones said understanding the downstream effects of BRAF mutation could allow for drug intervention to prevent this cascade of DNA changes from happening altogether. Ultimately, this might prevent the colon cancer from developing.

"The next phase is to look at how the changes in BRAF cause this cascade leading to cancer," he said. "This was a huge step in the right direction that could have clinical relevance for patients in a meaningful way."

The research was published in the scientific journal *PLOS ONE*. The research was supported by grant Nos. PO1CA073992 and R01CA116468 from the National Cancer Institute, a part of the National Institutes of Health.

Provided by Oklahoma Medical Research Foundation

Citation: New colon cancer finding could lead to earlier diagnosis—and better outcomes (2018, March 28) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-03-colon-cancer-earlier-diagnosisand-outcomes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.