

30-day ER revisit predicts poor outcomes in elderly

March 8 2018



(HealthDay)—A revisit to the emergency department within 30 days of a



previous visit predicts poor outcomes in elderly adults, according to a study published online Feb. 28 in the *Journal of the American Geriatrics Society*.

Jelle de Gelder, M.D., from the Leiden University Medical Center in the Netherlands, and colleagues assessed predictors of <u>emergency</u> department revisits and the association between revisits and 90-day functional decline or mortality among 1,093 <u>older adults</u> (median age 79 years) discharged from the emergency departments of one academic and two regional Dutch hospitals.

The researchers found that 114 participants (10.4 percent) had an emergency department revisit within 30 days of discharge. Independent predictors of a 30-day revisit included age (hazard ratio [HR], 0.96), male sex (HR, 1.61), polypharmacy (HR, 2.06), and cognitive impairment (HR, 1.71). To predict a revisit, the area under the receiver operating characteristic curve was 0.65. Individuals with a revisit were at higher risk (odds ratio, 1.99) of functional decline or mortality in propensity-score-matched analysis.

"An early emergency department revisit is a strong new predictor of adverse outcomes in older <u>adults</u>," the authors write.

More information: <u>Abstract/Full Text</u>

Copyright © 2018 <u>HealthDay</u>. All rights reserved.

Citation: 30-day ER revisit predicts poor outcomes in elderly (2018, March 8) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2018-03-day-er-revisit-poor-outcomes.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.