

Diaphragmatic breathing tied to less belching in GERD

March 5 2018



(HealthDay)—Diaphragmatic breathing can reduce belching and proton

pump inhibitor (PPI)-refractory gastroesophageal reflux, according to a study published in the March issue of *Clinical Gastroenterology and Hepatology*.

Andrew Ming-Liang Ong, M.B.Ch.B., from Singapore General Hospital, and colleagues investigated whether diaphragmatic [breathing](#) therapy was effective in reducing belching and PPI-refractory [gastroesophageal reflux](#) disease (GERD) symptoms among 36 consecutive [patients](#) seen from April 2015 through October 2016. Fifteen patients were placed on a standardized diaphragmatic breathing exercise protocol, and 21 were placed on a wait list and eventually received the breathing therapy.

The researchers found that 60 percent of patients in the treatment group and none of the control subjects achieved ≥ 50 percent reduction in belching visual analogue scale score. In the breathing treatment group, 80 percent of patients reduced belching frequency versus 19 percent in control subjects. Furthermore, the breathing treatment was associated with significant reduction in symptoms of GERD and a significant increase in quality of life scores. Four months post-treatment, these changes were sustained.

"Diaphragmatic breathing exercise training can be a useful adjunct to current pharmacologic options in treating these patients," the authors write.

More information: [Abstract/Full Text](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Diaphragmatic breathing tied to less belching in GERD (2018, March 5) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-03-diaphragmatic-tied-belching-gerd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.