

# Diuretic therapy for extremely preterm infants does not alleviate respiratory problems

March 27 2018

---

Diuretic therapy—commonly given to extremely preterm infants to help them overcome respiratory problems—appears to offer no benefit for this purpose, according to an analysis by researchers at the National Institutes of Health. Surprisingly, infants in the study who received diuretic therapy were more likely to require respiratory support, compared to extremely preterm infants with similar respiratory problems who did not receive the therapy. The study is published in the *Journal of Pediatrics*.

[Diuretic medicines](#) prompt the kidneys to make more urine. The [therapy](#) is commonly given to preterm [infants](#) to help drain fluid from the lungs, but there is little research evidence to support the practice.

Researchers at NIH's Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) and National Heart, Lung, and Blood Institute (NHLBI) analyzed data collected as part of the Prematurity and Respiratory Outcomes Program study, an effort to understand respiratory difficulties in extremely premature infants during the first year of life. The study included 835 infants born between 23 and 28 weeks at 13 U.S. neonatal intensive care units. An infant is considered [full term](#) at 39 weeks.

Researchers analyzed the infants' daily medication and respiratory support records from birth through 34 weeks. They found that 90

percent of the infants on diuretics needed supplemental oxygen, compared to 59 percent of infants who did not receive the therapy. On the day prior to the first diuretic treatment (14 days of life), 56 percent of the diuretic group were sicker and on ventilator support, compared to 11 percent of the non-diuretic group.

"Prescribing patterns for diuretics vary among hospitals," said Anne Zajicek, M.D., Pharm. D., the study's senior author. Dr. Zajicek was at NICHD when the analysis was conducted and is now deputy director of the NIH Office of Clinical Research. "We set out to explore whether this therapy helps very premature infants, but we found that it may be ineffective, which was contrary to what we expected."

The authors called for more research to better understand diuretic treatments. NICHD currently is funding a study looking at the effectiveness of furosemide, the most commonly used diuretic, for treating bronchopulmonary dysplasia, a lung condition affecting infants who have undergone ventilator therapy or oxygen therapy.

"The study reinforces growing concerns about the risk and possible benefits of diuretics, which are one of the most commonly prescribed classes of medications in the [neonatal intensive care](#) unit," said Carol Blaisdell, M.D., the study's principal investigator. Dr. Blaisdell was at NHLBI when the analysis was conducted and is now a senior program officer in the Environmental Influences and Child Health Outcome program within the NIH Office of the Director.

Provided by National Institutes of Health

Citation: Diuretic therapy for extremely preterm infants does not alleviate respiratory problems (2018, March 27) retrieved 4 May 2024 from <https://medicalxpress.com/news/2018-03-diuretic-therapy-extremely-preterm-infants.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.