


Higher positive cardiomyopathy genetic detection rates for whites

March 5 2018



(HealthDay)—Compared with underrepresented minorities, white

individuals have significantly higher positive detection rates and lower rates of inconclusive results on cardiomyopathy genetic testing, according to a study published online Feb. 28 in *JAMA Cardiology*.

Latrice G. Landry, Ph.D., from the U.S. Food and Drug Administration in Silver Spring, Md., and Heidi L. Rehm, Ph.D., from Brigham and Women's Hospital in Boston, conducted a cross-sectional analysis of the genetic panel [test](#) results for 5,729 probands with a suspected diagnosis or family history of cardiomyopathy who had been referred for testing. The results were stratified into three categories: white (79.2 percent), Asian (6.1 percent), and underrepresented minorities (14.7 percent).

The researchers found that positive detection occurred in 29 percent of white individuals compared with 18.4 and 25 percent of underrepresented minorities and Asians, respectively (P

"This clear disparity warrants further study to understand the gaps in usefulness, which may derive from a lack of clinical testing and research in underrepresented minority populations, in the hopes of improving genetic testing outcomes for [cardiomyopathy](#) in nonwhite groups," the authors write.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

[Editorial \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Higher positive cardiomyopathy genetic detection rates for whites (2018, March 5) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2018-03-higher-positive-cardiomyopathy-genetic-whites.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.