


Intrauterine adhesions rarely occur post induced abortion

March 23 2018



(HealthDay)—Intrauterine adhesions (IUA) following an induced

abortion are rare, but risk increases when surgical evacuation is involved, according to a study published online March 13 in *BJOG: An International Journal of Obstetrics & Gynaecology*.

Maarit Mentula, M.D., from the University of Helsinki, and colleagues retrospectively assessed data from a nationwide registry to identify women who had an induced [abortion](#) in Finland between 2000 and 2008, as well as matched controls.

The researchers identified a total of 12 (1.5 per 10,000) IUA diagnoses from 79,960 eligible induced abortions. The rate of IUA was 1.5 per 10,000 medically-induced abortions and 2.0 cases per 10,000 surgically-induced abortions ($P = 0.19$). In a [subgroup analysis](#), [surgical treatment](#) of the remaining products of conception following abortion significantly increased the risk of IUA (odds ratio, 5.50; $P = 0.012$).

"These results suggest that trauma to a recently pregnant uterus is an important risk factor for IUA," the authors write.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Intrauterine adhesions rarely occur post induced abortion (2018, March 23) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-03-intrauterine-adhesions-rarely-abortion.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--