


# LSD blurs boundaries between the experience of self and other

March 19 2018




Credit: CC0 Public Domain

LSD reduces the borders between the experience of the self and others, and thereby affects social interactions. Researchers at the University of Zurich have now found that the serotonin 2A receptor in the human brain is critically involved in these intertwined psychological mechanisms . This knowledge could contribute to new therapies for psychiatric disorders such as schizophrenia or depression.

Virtually all [mental health disorders](#) come with difficulties in interpersonal relations that in the long run negatively affect the progression of the disease. The associated health and social restrictions can only be marginally improved by current forms of therapy. One of the reasons for this is that there has been very little research into the basic neurobiological principles and in particular the neurochemical mechanisms of these kinds of disorders. A further symptom of various [psychiatric disorders](#) is distortions of self-experience. People suffering from mental disorders often show either an inflated or weakened sense of self.

Researchers at the University Hospital of Psychiatry Zurich have now investigated the links between changes in the sense of self and changes in [social interaction](#), and the pharmacological mechanisms that play a role in these processes.

"LSD blurs the boundaries between one's own self and others during a social interactions," explains Katrin Preller. She led the research team of the Neuropsychology and Brain Imaging group together with Prof. Franz Vollenweider and cooperated with the Max Planck Institute of Psychiatry in Munich on the study. While lying in an MRI scanner, study participants communicated with a virtual avatar by means of eye movements after having been administered a placebo, LSD, or LSD in combination with ketanserin.



LSD reduced activity in the posterior cingulate cortex and the temporal cortex, brain areas important for establishing one's sense of self. Credit: Preller et al., *JNeurosci* (2018)

### Changes in social interaction

"This allowed us to show that brain regions which are important for distinguishing between self and others were less active under the influence of LSD," says Preller. "And this also changed social interactions." The researchers were also able to show that the LSD-induced changes were blocked by ketanserin indicating that the serotonin 2A receptor (5-HT<sub>2A</sub> receptor) plays a critical role in this mechanism.

## Approaches for new drugs

These findings demonstrate that self-experience and social interaction are closely linked. Varying impairments of these intertwined processes could be the result of an impaired transfer of information mediated by the 5-HT<sub>2A</sub> receptor system. This could be important for the development of new pharmacological therapies. For example, blocking this receptor in patients suffering from an incoherent sense of self such as schizophrenia could improve their symptoms as well as their social abilities. On the other hand, stimulating this receptor could help patients who suffer from an increased self-focus, as is the case with depression, for example.

**More information:** *Journal of Neuroscience*, [DOI: 10.1523/JNEUROSCI.1939-17.2018](https://doi.org/10.1523/JNEUROSCI.1939-17.2018)

Provided by University of Zurich

Citation: LSD blurs boundaries between the experience of self and other (2018, March 19)  
retrieved 21 July 2024 from <https://medicalxpress.com/news/2018-03-ld-blurs-boundaries.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--