


Mice increase wheel running, activation of key brain regions two hours before methamphetamine availability

March 12 2018



Animals live in the home cage that is connected to the nebulizing chamber by a tunnel. The orbitofrontal cortex (OFC), lateral septum (LS) and dorsomedial hypothalamus (DMH) are activated in anticipation of access to the nebulizing chamber when methamphetamine, but not water, is available. Credit: Juárez-Portilla et al., *eNeuro* (2018)



The brain regions activated in anticipation of methamphetamine are identified in a noninvasive study of male mice published in *eNeuro*.

Rae Silver and colleagues investigated behavioral and <u>brain changes</u> in response to regularly scheduled access to <u>methamphetamine</u>, one of the most commonly abused drugs worldwide. Mice were housed in a cage with food, water and a running wheel connected via tunnel to a separate chamber where either nebulized methamphetamine or water was available. The door to the nebulizing chamber was open for one hour at the same two time periods (early, late) each day. This experimental design avoids complications associated with injections and surgery and better represents how humans use the drug.

The researchers found that mice in the methamphetamine group, but not the water group, increased their running on the wheel two hours before the door to the nebulizing <u>chamber</u> was open. The activated <u>brain</u> <u>regions</u> included those implicated in reward processing, food anticipation and craving: the orbitofrontal cortex, the dorsomedial hypothalamus and lateral septum.

This anticipation effect was stronger early in the day, a time of day finding consistent with a previous report in humans that found admission of overdose patients to an emergency room peaked at about 6:30 p.m.

More information: Brain activity during methamphetamine anticipation in a non-invasive self-administration paradigm in mice, *eNeuro*, <u>DOI: 10.1523/ENEURO.0433-17.2018</u>

Provided by Society for Neuroscience

Citation: Mice increase wheel running, activation of key brain regions two hours before



methamphetamine availability (2018, March 12) retrieved 8 May 2024 from https://medicalxpress.com/news/2018-03-mice-wheel-key-brain-regions.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.