

## Mono-unsaturated fats from plants, not animals may reduce risk of death from heart disease and other causes

## March 21 2018

Diets rich in mono-unsaturated fatty acids from plants were associated with a lower risk of dying from heart disease or other causes compared to diets rich in mono-unsaturated fats from animals, which were linked to a higher risk of death from heart disease or other causes, according to preliminary research presented at the American Heart Association's Epidemiology and Prevention / Lifestyle and Cardiometabolic Health Scientific Sessions 2018, a premier global exchange of the latest advances in population based cardiovascular science for researchers and clinicians.

"Our results emphasize the importance of the source and quantity of mono-unsaturated fatty acids in the diet - we should eat more mono-unsaturated fatty acids from plant sources and less mono-unsaturated fatty acids from animal sources," said Marta Guasch-Ferré, Ph.D., a research associate and one of the lead authors of this study along with Geng Zong, Ph.D., a research fellow. Both are at the Harvard School T.H. Chan of Public Health in Boston.

Mono-unsaturated fats are usually liquid at room temperature and solidify when refrigerated. Sources of plant-based mono-unsaturated fats include olive and other vegetable oils, avocados and many nuts and seeds. Sources of animal-based mono-unsaturated fats include full-fat dairy products, eggs, poultry, red meats and fish.



To assess the impact of mono-unsaturated fatty acids consumption on death from cardiovascular disease and other causes, researchers used data from 63,412 women from the Nurses' Health Study and 29,966 men from the Health Professionals Follow-Up Study. Both studies used detailed food-frequency questionnaires administered every four years to evaluate the composition of the participants' diets. This type of observational study can identify a trend among the participants but cannot prove cause and effect.

During an average 22 years of follow-up, there were 20,672 deaths among participants, 4,588 of them from heart disease. Analyzing the diet information, the researchers found:

- Participants with a higher intake of mono-unsaturated fatty acids from plants had a 16 percent lower risk of death from any cause compared to those with lower intakes.
- Participants with a higher intake of mono-unsaturated fatty acids from animals had a 21 percent higher risk of death from any cause.
- Replacing saturated fats, refined carbohydrates (like simple sugars) or trans fats with an equal number of calories (2 percent 5 percent of the total) from mono-unsaturated fatty acids from plants might lower the risk of heart disease deaths and death from any cause between 10 percent and 15 percent.
- Replacing mono-unsaturated fatty acids from animals with an equal amount of calories (5 percent of the total) of monounsaturated fatty acids from animals might lower the risk of heart disease deaths and deaths from any cause between 24 percent to 26 percent.

In the study, the risks were adjusted to account for several known factors that could influence the risk of death, including ethnicity; smoking status; intake of alcohol, fruits and vegetables and total calories; family



history of chronic diseases; physical activity; body mass index; and heart disease risk factors when participants enrolled. The results should be interpreted with caution because the study relied on the participants' self-reporting what they are and because participants consuming higher amounts of plant-based foods may be more health conscious in general.

## Provided by American Heart Association

Citation: Mono-unsaturated fats from plants, not animals may reduce risk of death from heart disease and other causes (2018, March 21) retrieved 25 April 2024 from <a href="https://medicalxpress.com/news/2018-03-mono-unsaturated-fats-animals-death-heart.html">https://medicalxpress.com/news/2018-03-mono-unsaturated-fats-animals-death-heart.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.