

New study on nicotinic receptors and LT memory paves way for targeted dementia therapy

March 27 2018



Credit: CC0 Public Domain

A new University of Bristol study, which identifies how acetylcholine impacts learning and memory by acting at different receptors, could

prove significant in the drive to develop more targeted and effective therapies for dementia.

Currently, the main treatments for Alzheimer's disease are drugs that increase levels of [acetylcholine](#) in the brain. The University of Bristol paper, published today in *Cell Reports*, describes the role two main types of nicotinic receptor ($\alpha 7$ and $\alpha 4\beta 2$ [receptors](#)) play in [long-term memory](#) retrieval and encoding - and the impact of acetylcholine on these receptors.

Nicotinic receptors, named after their sensitivity to nicotine, are expressed in the human and rodent brain where they are naturally activated by the neurotransmitter acetylcholine. Activation of nicotinic receptors by acetylcholine in the frontal cortex of the brain has previously been shown to be essential for functions such as attention and working memory.

Professor of Cellular Neuroscience, Zafar Bashir, and PHD student, Marie Sabec, in the School of Physiology, Pharmacology and Neuroscience, have demonstrated for the first time that both types of nicotinic receptor are essential for long-term associative recognition in rats.

What's more, the study also found that each nicotinic receptor subtype was responsible for distinct aspects of memory. The initial encoding of information ("learning") was dependent on the $\alpha 7$ nicotinic receptor, and subsequent memory retrieval ("remembering") relied on the $\alpha 4\beta 2$ receptor.

Prof. Bashir commented on the findings:

"Learning is thought to rely on changes in the strength of communication between neurons. In this study we have shown that the $\alpha 7$ nicotinic

receptors enhance communication between the hippocampus and [frontal cortex](#) but $\alpha 4\beta 2$ nicotinic receptors decrease communication between these regions. Therefore, acetylcholine acting on different subtypes of nicotinic receptor in the frontal brain can enhance or depress neural communication for the learning and remembering of long-term memory, respectively.

"These findings could have significant implications for the way stimulation of acetylcholine is used to treat Alzheimer's. If drugs can be developed to target the individual [nicotinic receptors](#), responsible for different aspects of long-term memory, we could see much more targeted and effective therapies for dementia."

The paper 'Nicotinic acetylcholine receptors control encoding and retrieval of associative recognition memory through plasticity in the [medial prefrontal cortex](#)' by Sabec et al is published today (27 March 2018) in *Cell Reports*.

Provided by University of Bristol

Citation: New study on nicotinic receptors and LT memory paves way for targeted dementia therapy (2018, March 27) retrieved 24 June 2024 from <https://medicalxpress.com/news/2018-03-nicotinic-receptors-lt-memory-paves.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--