

Obesity surgery prevents severe chronic kidney disease and kidney failure

March 23 2018



Per-Arne Svensson, associate professor, Institute of Health and Care Sciences, Sahlgrenska Academy, Sweden Credit: Malin Arnesson

Patients that underwent weight-loss surgery ran a significantly lower risk of developing severe chronic kidney disease and kidney failure when

compared to conventionally treated patients, according to a study published in *International Journal of Obesity*.

Obesity is a dangerous condition accompanied by elevated blood pressure, cardiovascular diseases and type 2 diabetes, which in turn may lead to complications such as chronic kidney [disease](#). It is also an independent risk factor in developing chronic kidney disease. Different obesity treatment strategies include lifestyle interventions, dietary modification, pharmacological and surgical treatment.

Many recent studies of [patients](#) with obesity have shown that weight-loss (bariatric) surgery is currently the most efficient way to achieve and maintain significant long-term weight loss and prevent obesity-related complications such as type 2 diabetes. In the Swedish Obese Subjects (SOS) study conducted by Sahlgrenska Academy, researchers followed over 4000 patients with obesity for more than 20 years. Half of the patients received [weight-loss surgery](#), and the other half were treated with conventional non-surgical methods through primary health care.

The SOS researchers published their data in the *International Journal of Obesity*, reporting the incidence of advanced chronic kidney disease among patients in the SOS study. The results demonstrate that the patients that underwent weight-loss surgery ran a significantly lower risk of developing severe [chronic kidney disease](#) and [kidney failure](#) when compared to conventionally treated patients.

Patients with evidence of kidney damage (high levels of protein in the urine) at the start of the study benefited most from surgical treatment, indicating that surgery prevents progression of pre-existing kidney injury towards renal failure. This complements previous reports from the SOS group that showed that bariatric surgery prevented new onset kidney injury.

These [kidney](#)-specific findings add to a large body of evidence from the SOS study demonstrating that bariatric surgery reduces mortality, improves type 2 diabetes and prevents cardiovascular disease and cancer.

More information: A. Shulman et al, Incidence of end-stage renal disease following bariatric surgery in the Swedish Obese Subjects Study, *International Journal of Obesity* (2018). [DOI: 10.1038/s41366-018-0045-x](#)

Provided by University of Gothenburg

Citation: Obesity surgery prevents severe chronic kidney disease and kidney failure (2018, March 23) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-03-obesity-surgery-severe-chronic-kidney.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.