

# Sitting and physical inactivity may increase risk of urinary tract symptoms

March 21 2018

---

Prolonged sitting time and low physical activity levels were linked with the development of lower urinary tract symptoms (LUTS) in a *BJU International* study of 69,795 middle-aged Korean men.

A team led by researchers at the Kangbuk Samsung Hospital, in South Korea, found that the incidence rate of LUTS—which relate to urine storage and/or voiding disturbances—was 39 per 1000 person-years. (A person-year is the number of years of follow-up multiplied by the number of people in the study.)

"The results support the importance of both reducing sitting time and promoting physical activity for preventing LUTS," said lead author Dr. Heung Jae Park. "Further studies are still needed to examine the influence of sedentary behaviors on LUTS and its determinants," added senior author Dr. Seungho Ryu.

**More information:** Heung Jae Park et al, Sitting time, physical activity and the risk of lower urinary tract symptoms: a cohort study, *BJU International* (2018). [DOI: 10.1111/bju.14147](https://doi.org/10.1111/bju.14147)

Provided by Wiley

Citation: Sitting and physical inactivity may increase risk of urinary tract symptoms (2018, March 21) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-03-physical->

[inactivity-urinary-tract-symptoms.html](http://inactivity-urinary-tract-symptoms.html)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.