

Poor outcomes common with childhood anorexia nervosa

March 5 2018



(HealthDay)—Anorexia nervosa (AN) that develops before age 14 often

leads to unfavorable outcomes, according to a study published online Feb. 16 in the *International Journal of Eating Disorders*.

Beate Herpertz-Dahlmann, M.D., from University Clinics RWTH Aachen in Germany, and colleagues followed up with 52 former pediatric patients (mean age, 12.5 [SD 1.0]) diagnosed with early-onset AN who had been treated at German university hospitals. Each participant had a personal examination after an average of 7.5 years.

The researchers found that 41 percent of the participants had a good outcome, while 35 percent had intermediate outcomes, and 24 percent had poor outcomes. Twenty-eight percent of respondents met the DSM-IV criteria for a current non-eating disorder psychiatric disorder, and 64 percent met the criteria for a previous non-eating disorder psychiatric disorder. AN outcomes were strongly associated with mental-health-related quality of life and eating disorder-specific psychopathology. The only significant positive indicator of outcome was a higher weight at admission.

"Childhood AN is a serious disorder with an unfavorable course in many patients and high rates of chronicity and [psychiatric comorbidity](#) in young adulthood," the authors write. "Early detection and intervention are urgently needed."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Poor outcomes common with childhood anorexia nervosa (2018, March 5) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-03-poor-outcomes-common-childhood-anorexia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.