


From signal propagation to consciousness: New findings point to a potential connection

March 22 2018



Credit: CC0 Public Domain

Researchers at New York University have discovered a novel mechanism through which information can be effectively transmitted across many areas in the brain—a finding that offers a potentially new way of

understanding how consciousness arises.

The study appears in the journal *Neuron*.

"For us to navigate through everyday situations, it is critical that [electrical signals](#) from our senses stimulated by the external world are able to reach relevant brain [areas](#)," explains Xiao-Jing Wang, Global Professor of Neural Science at NYU and the paper's senior author. "Our brain, however, is a very complicated system, with billions of [neurons](#) organized in an interconnected network with more than a hundred areas. It is therefore not easy to understand how signals may travel from area to area in an efficient manner."

Brain areas tend to be organized in a hierarchy, ranging from "lower" sensory areas to "higher" cognitive areas. These areas have excitatory and [inhibitory neurons](#), which either stimulate or suppress activity in other neurons.

Understanding how [neural signals](#) are transmitted across this hierarchy, the researchers note, is still a fundamental challenge in neuroscience and served as the focal point for the *Neuron* study.

Previous computer modeling of signal transmission across areas in the brain did not take into account the complexity of brain's area-to-area connections. By contrast, thanks to the recent advances in the field of "Connectomics" dedicated to quantitatively analyzing the brain's connectivity, the NYU scientists were able to build models incorporating anatomical connectivity data of macaque monkeys.

Here they found that signal transmission in a large-scale model of the primate brain is robust under the condition in which area-to-area connections exhibit a "balance" between excitation and inhibition. Specifically, stimulation provided by the excitatory neurons allows for

signals to be transmitted, whereas suppression from inhibitory neurons makes sure the signal activity does not grow out of control.

"Unexpectedly, our model reveals that only when the signal is strong enough, above a threshold level, the signal reaches a large set of areas of the brain region called the prefrontal cortex, which plays a critical role in high-level cognition," said Madhura Joglekar, the paper's first author who conducted the research as a postdoctoral fellow in Wang's lab and is now an instructor at NYU's Courant Institute of Mathematical Sciences.

Notably, the global activation pattern resembles the ones previously found in the human [brain](#) while consciously perceiving sensory information—a parallel suggesting a potential connection between the proposed "balanced" large-scale neural circuit mechanism for [signal transmission](#) and understanding how conscious information processing is achieved.

Provided by New York University

Citation: From signal propagation to consciousness: New findings point to a potential connection (2018, March 22) retrieved 1 May 2024 from <https://medicalxpress.com/news/2018-03-propagation-consciousness-potential.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--