


Racial disparities in HIV control persist despite equal access to care

March 21 2018



Credit: Mary Ann Liebert, Inc., publishers

Researchers report that racial disparities in HIV control (viral load) exist even when patients have equal access to care, as shown in a study of black and white HIV-infected patients treated in the Veterans Administration (VA) health system. The study, which identified specific factors that contributed to these differences, is published in *AIDS Patient Care and STDs*.

The article "Patient and Facility Correlates of Racial Differences in Viral Control for Black and White Veterans with HIV Infection in the Veterans Administration" was coauthored by Mary Vaughan Sarrazin, PhD, Iowa City VA Medical Center and University of Iowa (Iowa City) and colleagues from VA Palo Alto Health Care System (CA), Stanford University (CA), ENRM Veterans Affairs Medical Center (Bedford, MA), and Boston University School of Public Health (MA). The researchers determined that more than half of the disparity in viral control between black and white veterans could be attributed to differences in adherence to [combination antiretroviral therapy](#) (cART).

"This is an especially critical issue given the widening disparity in incidence of new HIV infections between African Americans and other racial/ethnic groups," says Editor-in-Chief Jeffrey Laurence, MD, Professor of Medicine, Weill Cornell Medical College, New York, NY.

More information: *AIDS Patient Care and STDs* (2018). [DOI: 10.1089/apc.2017.0213](#)

Provided by Mary Ann Liebert, Inc

Citation: Racial disparities in HIV control persist despite equal access to care (2018, March 21) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-03-racial-disparities-hiv-persist-equal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.