

# Women report fewer adverse side effects from partial or reduced breast radiotherapy

March 23 2018

---

The average number of moderate or marked side-effects reported by breast cancer patients is lower if they are treated with radiotherapy to part of the breast or a reduced dose to the whole breast, rather than with standard radiotherapy to the whole breast, according to new findings presented at the 11th European Breast Cancer Conference on Friday.

The IMPORT LOW study of 2016 women in 41 centres in the UK has already shown that partial [breast](#) and reduced dose [radiotherapy](#) was as effective as whole breast radiotherapy in controlling the cancer at five years, and women in the partial breast and reduced dose groups reported fewer side-effects, including less change in the appearance of the breast.

These latest results, which focus predominantly on side-effects affecting the breast and also body image reported during the five years following radiotherapy, show that over half of [patients](#) in the study did not report moderate or marked side-effects at any point and that most side-effects reduced over time. The number of side-effects reported per person were fewer in the partial breast and reduced dose groups compared with the whole breast radiotherapy group.

Dr Indrani Bhattacharya, a clinical research fellow at the Institute of Cancer Research's Clinical Trials and Statistics Unit (London, UK), from where the IMPORT LOW trial was coordinated, told the conference: "The findings from this study are reassuring for women who are offered either whole breast or partial breast radiotherapy using this technique of radiotherapy, which is simple to deliver and already

available in centres worldwide.

"This new information will help doctors discuss the risks and benefits of this type of radiotherapy with patients and may improve shared decision-making, as well as enabling them to tailor treatment for individual patients."

The IMPORT LOW study randomised women to receive standard radiotherapy at a dose of 40 Gy to the whole breast (the control group), or 36 Gy to the whole breast and 40 Gy to the part of the breast that contained the original tumour (reduced dose group), or 40 Gy only to the site of the original tumour (partial breast group). The radiotherapy was given using hypofractionated intensity modulated radiotherapy (IMRT) - a technique that can deliver an even dose of radiation, minimising the chances of hotspots of unwanted high doses and reducing the cosmetic problems that can occur after breast radiotherapy.

More than half of the women (1265) took part in the sub-study that looked at patient-reported outcomes (PROMs). At the start of the study, after six months and one, two and five years after radiotherapy the women were asked about adverse side-effects, such as a hardening of tissue, pain, over-sensitivity of the treated area and build-up of fluid. Most of the side-effects reported by the women related to changes affecting the breast, and the commonest reported side-effect was "overall change in breast appearance".

The researchers found that women were more likely to report adverse side-effects if they were younger, had larger breasts, had a larger volume of breast tissue removed at surgery, if the cancer had spread to any of their lymph nodes, and if, at the start of the study, they were feeling anxious or depressed.

"Now that we can identify these patients who are at higher risk of

reporting side-effects, this knowledge can be discussed with patients, may modify treatment and enable doctors to put in place more personalised and frequent monitoring if necessary. For example, patients with higher levels of anxiety and depression could be offered psychosocial support at the start of treatment, although we do not know from this study whether this would reduce the reporting of side effects," said Dr Bhattacharya.

"We have previously reported that partial breast radiotherapy is as effective as whole breast radiotherapy with similar or fewer side-effects over five years and can be safely delivered in a selected group of patients with low-risk breast cancers. This new analysis shows that patients requiring whole breast radiotherapy can be reassured about the low risk of side-effects affecting the breast and body image. The technique of radiotherapy used in IMPORT LOW is easy to implement and deliver as the equipment and expertise is available in all centres worldwide."

The IMPORT LOW trial is the largest randomised trial to study the use of IMRT to deliver partial or low dose breast radiotherapy compared to whole breast radiotherapy, and has provided the largest complete set of PROMs data for several points in time over five years.

"PROMs are very important for both women and clinicians as they describe side-effects from the patients' perspective and allow us to see the long-term effects of treatment as recorded by the patient. We know from this study and other studies that patients report more side-effects compared to doctors. This means that if we did not assess side-effects using PROMs, then the impact of the treatment-related side-effects on the patient may be underestimated," said Dr Bhattacharya.

Co-chair of the European Breast Cancer Conference, Professor Isabel Rubio, director of the Breast Surgical Unit at Clinica Universidad de Navarra (Spain), who was not involved with the research, commented:

"Patient-reported outcomes are among the most important factors that we have added to assessments of outcomes in breast cancer in recent years. PROMs can improve relationships between physicians and patients, and enhance shared decision-making; this strategy will impact on the final outcomes in the management of [breast cancer patients](#). These latest results from the IMPORT LOW trial provide important and reassuring information for clinicians and patients about side effects after partial or reduced dose radiotherapy, and will help them to make the best choices in each individual case."

**More information:** Abstract no: 3 (Best abstract), "A longitudinal analysis of patient reported outcomes over 5 years in the IMPORT LOW (partial breast radiotherapy after breast conservation surgery) phase III randomised controlled trial", Friday, closing plenary session

Provided by ECCO-the European CanCer Organisation

Citation: Women report fewer adverse side effects from partial or reduced breast radiotherapy (2018, March 23) retrieved 15 May 2024 from <https://medicalxpress.com/news/2018-03-women-adverse-side-effects-partial.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--