

# Attention deficit disorders could stem from impaired brain coordination

April 5 2018

---



The Case Western Reserve University School of Medicine. Credit: CWRU School of Medicine

Researchers from Case Western Reserve University School of Medicine and colleagues have discovered how two brain regions work together to

maintain attention, and how discordance between the regions could lead to attention deficit disorders, including schizophrenia, bipolar disorder, and major depression.

People with attention deficits have difficulty focusing and often display compulsive behavior. The new study suggests these symptoms could be due to dysfunction in a gene—*ErbB4*—that helps different [brain regions](#) communicate. The gene is a known risk factor for psychiatric disorders, and is required to maintain healthy neurotransmitter levels in the brain.

In a study published in the current issue of *Neuron*, researchers showed mice lacking *ErbB4* activity in specific brain regions performed poorly on timed attention tasks. The mice struggled to pay attention and remember visual cues associated with food. Neuroscientists describe the kind of thought-driven attention required for the tasks as "top-down attention." Top-down attention is goal-oriented, and related to focus. People who lack efficient top-down attention are at a higher risk for attention deficit hyperactivity disorder (ADHD). The study is the first to connect *ErbB4* to top-down attention.

"The results reveal a mechanism for top-down attention, which could go wrong in attention disorders," says corresponding author Lin Mei, PhD, professor and chair of the department of neurosciences at Case Western Reserve University School of Medicine. "And since *ErbB4* is a risk factor for schizophrenia, bipolar disorder, and major depression, the results provide insights into mechanisms of these disorders."

When the researchers attached probes to the mice to measure brain activity, they found mice without *ErbB4* had brain regions that were acting independently, rather than together in synchrony. In particular, the researchers studied the prefrontal cortex—normally associated with decision-making—and the hippocampus—a region that supports memory. These two regions coordinate for a variety of brain tasks,

including memory and attention. "We found top-down attention, previously thought to be controlled by the prefrontal cortex, also involves the hippocampus in a manner where the two regions are highly synchronized when attention is high," says Mei. "Our findings give importance to synchrony between the prefrontal cortex and hippocampus in top-down attention and open up the possibility that attention deficit disorders, like ADHD, might involve impairments in the synchrony between these two regions."

According to the new study, *ErbB4* coordinates a cascade of brain signals that "bridge" the two regions. *ErbB4* itself encodes a receptor found on the surface of brain cells. The study found that when a protein (neuregulin-1) attaches to the *ErbB4* receptor, it triggers a chain reaction that ultimately determines neurotransmitter levels in the prefrontal cortex and hippocampus. Without *ErbB4*, neurotransmitter levels go awry. The researchers discovered mice lacking *ErbB4* have low levels of a particular neurotransmitter—GABA, or gamma-aminobutyric acid—in their brain. Low GABA levels can lead to impaired top-down attention in the prefrontal cortex, and impairs how the [prefrontal cortex](#) can efficiently coordinate with the hippocampus. The researchers concluded that *ErbB4* helps link the two brain regions to maintain attention.

The study used a novel mouse model to study brain functions. By using genetic and chemical techniques, Mei's team can specifically inhibit *ErbB4* in a specific brain region. "We generated a mutant mouse that enables us to inhibit *ErbB4* activity whenever and wherever we want, thus allowing temporal and spatial control of *ErbB4* activity," says Mei. "This positions us to understand how different brain regions and their neurotransmitter activity regulate various brain functions." The researchers are planning to use the novel mouse model to study how *ErbB4* may coordinate [brain](#) activities, in an effort to learn more about mechanisms behind [attention](#) deficit [disorders](#).

**More information:** *Neuron* (2018). [DOI: 10.1016/j.neuron.2018.03.018](https://doi.org/10.1016/j.neuron.2018.03.018)

Provided by Case Western Reserve University

Citation: Attention deficit disorders could stem from impaired brain coordination (2018, April 5) retrieved 26 March 2023 from <https://medicalxpress.com/news/2018-04-attention-deficit-disorders-stem-impaired.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.