


Where the brain detects changes in natural sounds

April 9 2018



A. We presented natural textures which change their statistics at random times to a variable degree (depicted: 3 s). The change in statistics leads to a distributed change in spectro-temporal properties. The present example shows a transition from bubbling to a linear mixture between bubbling and rain, with an intermediate mixing coefficient of 0.3. Level is relative to the maximal overall level. B. Texture sounds were provided via headphones while simultaneously recording whole-head EEG signals from 64 channels. In the active paradigm, listeners were instructed to report whether they heard a change by pressing a button after the termination of the sound and otherwise not to respond. Instead, in the passive variant they were just asked to listen to the stimuli. Credit: Górska et al., *eNeuro* (2018)

Electrical activity in a region of the parietal cortex underlies the

detection of a transition between two complex sounds, finds a study of human participants published in *eNeuro*. The research provides insight into how the brain tunes into relevant changes in the environment to optimize behavior.

The world is filled with a multitude of ever-changing sound. While some of these sounds signal important information about the environment, such as the presence of danger, many others are not so useful. Detecting changes in a natural soundscape is essential for identifying which sounds should be attended to.

Bernhard Englitz and colleagues presented participants with random changes (or no change) between two acoustic textures resembling rain, applause or bubbling water through headphones while their brain activity was recorded using electroencephalography. The researchers found that the longer participants were exposed to the first texture, the faster their reaction time and ability to identify the changes. Activity in the precuneus, a part of the [parietal cortex](#) near the occipital lobe, appears to be the source of these effects, which were stronger when participants were prompted to indicate whether or not they noticed a change, but also present in passive listening conditions.

More information: Evidence integration in natural acoustic textures during active and passive listening, *eNeuro*, [DOI: 10.1523/ENEURO.0090-18.2018](#)

Provided by Society for Neuroscience

Citation: Where the brain detects changes in natural sounds (2018, April 9) retrieved 10 May 2024 from <https://medicalxpress.com/news/2018-04-brain-natural.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.