

Heading—not collisions—cognitively impairs players

April 24 2018



Credit: CC0 Public Domain

Worse cognitive function in soccer players stems mainly from frequent ball heading rather than unintentional head impacts due to collisions, researchers at Albert Einstein College of Medicine have found. The

findings suggest that efforts to reduce long-term brain injuries may be focusing too narrowly on preventing accidental head collisions. The study published online today in the *Frontiers in Neurology*.

"Unintentional head impacts are generally considered the most common cause of diagnosed concussions in soccer, so it's understandable that current prevention efforts aim at minimizing those collisions," said study leader, Michael Lipton, M.D., Ph.D., F.A.C.R., professor of radiology and of psychiatry and behavioral sciences at Einstein and medical director of MRI Services at Montefiore. "But intentional head impacts—that is, soccer ball heading—are not benign. We showed in a previous study that frequent heading is an underappreciated cause of concussion symptoms. And now we've found that heading appears to alter cognitive function as well, at least temporarily."

While heading has previously been associated with transient cognitive problems, the Einstein study is the first to compare the cognitive effects of heading to unintentional head impacts such as collisions. Three hundred and eight amateur soccer players in New York City filled out questionnaires detailing their recent (previous two weeks) soccer activity, including heading and unintentional head impacts. Participants also completed neuropsychological tests of verbal learning, verbal memory, psychomotor speed, attention and working memory. The players ranged in age from 18 to 55, and 78 percent were male.

Players headed soccer balls an average of 45 times during the two weeks covered by the questionnaire. During that time, about one-third of the players suffered at least one unintentional head [impact](#) (e.g., kicks to the head or head-to-head, head-to-ground, or head-to-goalpost collisions).

Players who reported the most headings had the poorest performance on psychomotor speed and attention tasks, which are areas of functioning known to be affected by [brain injury](#). Heading frequency also correlated

with poorer performance on the working memory task, although the association was of borderline significance. In contrast, unintentional [head](#) impacts were not related to any aspect of [cognitive performance](#).

The changes in cognitive function did not cause overt clinical impairment, the Einstein team reported. "However, we're concerned that subtle, even transient reductions in neuropsychological function from heading could translate to microstructural changes in the brain that then lead to persistently impaired function. We need a much longer-term follow-up study of more soccer players to fully address this question," said Dr. Lipton.

In the meantime, soccer players should consider reducing heading during practice and [soccer](#) games, Dr. Lipton advises. "Heading is a potential cause of brain injury," he says, "and since it's under control of the player, its consequences can be prevented."

The paper is titled, "Heading Frequency is More Strongly Related to Cognitive Performance than Unintentional Head Impacts in Amateur Soccer Players."

Provided by Albert Einstein College of Medicine

Citation: Heading—not collisions—cognitively impairs players (2018, April 24) retrieved 13 March 2024 from

<https://medicalxpress.com/news/2018-04-headingnot-collisionscognitively-impairs-players.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--