

Smoking may increase heart failure risk among African-Americans

April 16 2018

African Americans who smoke appear to be at far greater risk of developing heart failure than those who never smoked, or those who quit, according to new research in the American Heart Association's journal *Circulation*.

"Previous research has focused on [smoking](#) and atherosclerosis, or hardening of the arteries, but not enough attention has been given to the other bad effects of smoking on the [heart](#)," said Michael E. Hall, M.D., M.S., a cardiologist at the University of Mississippi Medical Center in Jackson and senior study author. "With increasing rates of heart failure, particularly among African Americans, we wanted to look at the link between smoking and heart failure."

Patients with heart failure are unable to pump enough blood and oxygen to their bodies to remain healthy. According to the American Heart Association, 1 of every 5 Americans over the age of 40 is expected to develop heart failure in their lifetime—and that number is growing.

The study included 4,129 participants in the Jackson Heart Study with a median follow up of 8 years. At enrollment, none of the participants (average age 54) had heart failure or hardening of the arteries, which can lead to heart failure. Among participants there were 2,884 people who never smoked, 503 who were current smokers and 742 who were former smokers. During the study period, there were 147 hospitalizations for heart failure.

The study found hospitalizations for heart failure were:

- Nearly three times more likely among current smokers;
- Three-and-a-half times more likely among current smokers who smoked a pack or more a day; and
- Twice as likely among those with a smoking history equivalent to smoking a pack a day for 15 years.

Researchers also found a link between current smoking and a larger [left ventricle](#), the heart's main pumping chamber, which showed early signs that the left ventricle was not working properly. Hall said these changes in the left ventricle's structure and function likely put a person at greater risk of developing heart failure.

Importantly, researchers did not find a link between former smokers and heart failure hospitalization or changes in the left ventricle.

The study took into account [high blood pressure](#), diabetes, body mass and other factors that might have biased results. Researchers said the association between smoking, heart failure hospitalizations and left ventricle changes remained even after also accounting for those participants who developed [coronary heart disease](#) during the study period.

Study limitations include the fact that the participants lived in only three counties in the Jackson, Mississippi metropolitan area, so findings may not be generalizable to African Americans living elsewhere.

"Still, the study clearly underscores the harms of smoking and the benefits of quitting," Hall said. As healthcare professionals, we would recommend that all patients quit smoking anyway, but the message should be made even more forcefully to patients at higher risk of [heart failure](#)."

More information: *Circulation* (2018). [DOI: 10.1161/CIRCULATIONAHA.117.031912](https://doi.org/10.1161/CIRCULATIONAHA.117.031912)

Provided by American Heart Association

Citation: Smoking may increase heart failure risk among African-Americans (2018, April 16)
retrieved 5 May 2024 from
<https://medicalxpress.com/news/2018-04-heart-failure-african-americans.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--