

Infant death study reveals dangerous sleep practices among babysitters, relatives, others

April 2 2018



Credit: CC0 Public Domain

Babies who died during their sleep while being watched by someone other than parents often had been placed in unsafe sleep positions, such as on their stomachs, or in unsafe locations, such as a couch, a new study

has found.

In response to the troubling findings, the researchers are urging parents to educate anyone who cares for their children about safe sleep practices and the risk of Sudden Infant Death Syndrome (SIDS), the leading cause of [death](#) among babies between 1 month and a year of age. "If someone else - a babysitter, relative, or [friend](#) - is taking care of your baby, please make sure that they know to place your baby on the back in a crib and without any bedding," said Rachel Moon, MD, of the University of Virginia School of Medicine, the UVA Children's Hospital and UVA's Child Health Research Center.

Reducing Risk of Sudden Infant Death Syndrome

The researchers reviewed more than 10,000 infant deaths and found that 1,375 occurred when a parent was not present. Among those 1,375 cases, they determined:

- Babies were less likely to be placed on their back, the sleep position recommended by the American Academy of Pediatrics, than when under parental care.
- Babies were more likely to be placed in sleep environments with objects that might prove hazardous. The AAP recommends that sleep spaces be free of toys and soft bedding, including blankets and sleep bumpers.
- 5 percent of licensed childcare providers placed the babies in a crib or bassinet, as recommended. Among babysitters, this number was 49.1 percent. Among relatives, the number was only 29.4 percent, and among friends it was 27.1 percent.
- 1 percent of childcare providers had placed infants in the recommended supine position (on the back), compared with only 38.4 percent of relatives, 38.6 percent of friends and 37.8 percent of babysitters.

- Deaths under the supervision of friends and relatives were most likely to occur while the babies were held or placed on an adult bed.

"A lot of relatives and friends may not be aware that babies are safest on their backs," said researcher Jeffrey Colvin, MD, of Children's Mercy Kansas City. "They may have raised children before we knew that this was safest."

Efforts to Prevent Sudden Infant Death

There was some encouraging news: While past studies found that many licensed childcare providers placed [infants](#) on their stomachs to sleep, the researchers noted that that same group was now the most likely non-parental supervisor to place [babies](#) in the recommended sleep position and in cribs. They suggest this may be a result of the educational efforts of the Safe to Sleep national campaign and changes in state regulations.

"It's always best to discuss where and how your baby should sleep," Moon said. "You can't make assumptions that the person with whom your baby is staying will know what is safest."

The researchers have published their findings in the *Journal of Pediatrics*. The study's authors were Elena Lagon, Moon and Colvin.

More information: Elena Lagon et al, Characteristics of Infant Deaths during Sleep While Under Nonparental Supervision, *The Journal of Pediatrics* (2018). [DOI: 10.1016/j.jpeds.2018.01.051](https://doi.org/10.1016/j.jpeds.2018.01.051)

Provided by University of Virginia Health System

Citation: Infant death study reveals dangerous sleep practices among babysitters, relatives, others (2018, April 2) retrieved 23 April 2024 from <https://medicalxpress.com/news/2018-04-infant-death-reveals-dangerous-babysitters.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.