

Late, but not too late—screening for olfactory dysfunction

April 20 2018



Credit: University of Duisburg-Essen, Essen, Germany

In a large population-based study of randomly selected participants in Germany, researchers found that participants aged 65-74 years with olfactory dysfunction showed impaired cognitive performance.

Interestingly, this strong association was not present in younger (55-64 years) or older (75-86 years) participants. Additionally, the effect was more present in women than men.

In neurodegenerative diseases like Parkinson's and Alzheimer's disease (AD), olfactory function is diminished. Further, [olfactory dysfunction](#) precedes the onset of cognitive impairment within AD, which highlights its potential as biomarker for early, preclinical diagnosis. Several studies suggest that olfactory dysfunction predicts progression from normal cognitive functioning to [mild cognitive impairment](#) and AD. There is little evidence for this association concerning different age stages and gender differences.

The Heinz Nixdorf Recall (Risk Factors, Evaluation of Coronary Calcium and Lifestyle) study is an observational, population-based, prospective study that examined 4,814 participants (baseline: 45-75 years; 50% men) in the metropolitan Ruhr Area. Participants returned for two examinations every five years. For this analysis, 2,640 participants from the third examination were divided into anosmics (lack of olfactory function), hyposmics (impaired olfactory function) and normosmics (normal olfactory function) according to their Sniffin' Sticks Screening test scores. To examine age- and gender-specific associations, stratified analyses for gender and three [age groups](#) were conducted. Men and women differed significantly in their olfactory function. Women had higher scores on the olfactory test. Particularly, middle-aged anosmic male and female participants showed worst

performance in several cognitive tests. No associations were seen in younger and older adults. There was a quantitative association in all age groups as anosmics performed worse than hyposmics and hyposmics performed worse than normosmics in all subtests.

The found association in the middle-aged group might occur because this age band between 65-74 years is critical for the onset of age-related cognitive and olfactory decline. Regarding the older age group, the association may be covered by other risk factors occurring in this age band like hypertension, diabetes or coronary artery disease. The pathology potentially causing olfactory and [cognitive impairment](#) mainly occurs after the age of 65. Thus, the young-aged groups are cognitively healthier and are less often anosmics. This might account for the missing associations in this age [group](#). Nevertheless, [cognitive performance](#) was reflected by the decrement in olfactory function in all age groups. Participants with worst olfactory function showed worst cognitive performance and vice versa. More distinct effects were found for women compared to men. General differences in olfactory function between men and women can be the cause of this result. At this point we cannot conclude clinical implications regarding gender.

This is the first study reporting on age-specific associations of olfactory function and cognitive performance in the general population. Testing olfactory [function](#) is an easy and inexpensive way to detect dysfunctions and can help to identify individuals at risk of cognitive decline. Assessing [olfactory function](#) may be an appropriate marker to detect persons at risk of cognitive decline, especially in a crucial age stage between 65 and 74 years.

More information: Sarah Tebrügge et al, Olfactory Function is Associated with Cognitive Performance: Results of the Heinz Nixdorf Recall Study, *Journal of Alzheimer's Disease* (2018). [DOI: 10.3233/JAD-170863](#)

Provided by IOS Press

Citation: Late, but not too late—screening for olfactory dysfunction (2018, April 20) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2018-04-late-latescreening-olfactory-dysfunction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.