

New research suggests possible link between sudden infant death syndrome and air pollution

April 20 2018



The study found evidence suggesting an association between SIDS and exposure to airborne pollutants. Credit: University of Birmingham

A study led by the University of Birmingham suggests a possible



association between exposure to certain pollutants and an increased risk of so-called 'cot death'.

The research, published in *BMJ Open* and carried out in collaboration with the University of Oulu in Finland and the Medical Research Council Unit The Gambia in Africa, examines the relationship between the effects of short-term variations in <u>air pollution</u> and the onset of cot death or Sudden Infant Death Syndrome (SIDS).

The study found evidence suggesting an association between SIDS and exposure to larger particulate matter (<u>airborne pollutants</u>) called PM10, as well as nitrous dioxide (NO2). Other pollutants were not found to be associated with SIDS. The researchers looked at levels of air pollution the day before a SIDS death and compared them to levels on a previous reference day. The study involved over 200 SIDS cases in the West Midlands, between 1996 and 2006.

The research found an increased risk for SIDS two days after exposure to NO2, while exposure to PM10 was shown to have an effect for up to five days after exposure.

Lead author Dr. Ian Litchfield, of the University of Birmingham's Institute of Applied Health Research, said: "Certain groups of the population are more vulnerable to ambient air pollution than others and children figure predominantly among them due to the fragility of their immune system and the ratio of their lung capacity to their size.

"Concern around the effects of air pollution continues to mount particularly within developing economies where it contributes to 3.3 million premature deaths worldwide per year, a figure estimated to double by 2050 if the issue remains unattended.

"However, government policies appear slow to react, for example, in the



UK pledges to cease sales of diesel and petrol cars do not come into effect until 2040.

"Our study has highlighted that more research needs to be carried out to better understand the effects of air pollution on child health, while government policy needs to bring about change with increasing urgency."

In terms of what parents can do to reduce the effects of exposure to pollution on their child, the study's authors draw attention to advice that suggests staying indoors on days with higher levels of pollution, as well as cleaning systems for air in the home such as air purifiers, though these can be expensive.

Francine Bates, Chief Executive of The Lullaby Trust which funded the University of Birmingham-led study, said: "The findings of the study give some cause for concern and parents should be aware that exposure of their child to air pollution carries health implications, including a possible increase in the risk of SIDS.

"However, further research is needed to understand more about the link between air pollution and SIDS and what can be done to reduce <u>exposure</u> of young children to <u>pollution</u>.

"For now we would advise parents to follow safer sleep guidelines which are known to reduce the risk of SIDS such as sleeping their baby on their back on a firm, flat surface, in a clear cot or Moses basket and not exposing them to second hand cigarette smoke."

More information: Is ambient air pollution associated with onset of sudden infant death syndrome? A case-crossover study in the UK. *BMJ Open*. DOI: 10.1136/bmjopen-2017-018341



Provided by University of Birmingham

Citation: New research suggests possible link between sudden infant death syndrome and air pollution (2018, April 20) retrieved 8 May 2024 from https://medicalxpress.com/news/2018-04-link-sudden-infant-death-syndrome.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.